

Evaluation of novel protease enzymes on growth performance and apparent ileal digestibility of amino acids in poultry: enzyme screening

by Walk, C.L., Pirgozliev, V., Juntunen, K., Paloheimo, M. and Ledoux, D.R.

Copyright, Publisher and Additional Information: This is the author accepted manuscript. The final published version (version of record) is available online via Oxford University Press

Please refer to any applicable terms of use of the publisher.

DOI: <https://doi.org/10.3382/ps/pey080>



Walk, C.L., Pirgozliev, V., Juntunen, K., Paloheimo, M. and Ledoux, D.R. 2018. Evaluation of novel protease enzymes on growth performance and apparent ileal digestibility of amino acids in poultry: enzyme screening. *Poultry Science*.

27 March 2018

1 **Evaluation of novel protease enzymes on growth performance and apparent ileal**
2 **digestibility of amino acids in poultry: enzyme screening**

3 C. L. Walk^{1*}, V. Pirgozliev², K. Juntunen³, M. Paloheimo³ and D. R. Ledoux⁴

4 ¹ AB Vista, Marlborough Wiltshire United Kingdom

5 ² Harper Adams University, Shropshire, United Kingdom

6 ³ Roal Oy, Rajamäki, Finland

7 ⁴ University of Missouri, Columbia United States

8 * corresponding author: carrie.walk@abvista.com

9 Running title: Novel proteases in poultry

10 **Abstract**

11 Three experiments were conducted to evaluate eight neutral serine and six acid
12 aspartic proteases on growth performance and apparent ileal amino acid digestibility (**AID**) of
13 poultts (Experiment 1) or chicks (Experiments 2 and 3). Two basal diets were formulated: a
14 nutrient adequate positive control (**PC**), formulated to meet or exceed the nutrient
15 requirements for poultts (Experiment 1) or chicks (Experiments 2 and 3) and a negative
16 control (**NC**) diet formulated to achieve 85% (Experiments 1 and 2) or 80% (Experiments 3)
17 of the requirement for protein and amino acids. Phytase was included in all diets to provide
18 500 FTU/kg and xylanase was included in all diets to provide 10,000 (Experiments 1 and 2)
19 or 16,000 (Experiments 3) BXU/kg. Proteases were supplemented in the NC diet at an
20 equivalent amount of enzyme protein to create 16 experimental diets. There were five
21 birds/pen and 10 replicate pens per treatment in each experiment. In experiment 1, birds fed
22 the PC diet gained more ($P < 0.05$) than birds fed the NC. There were no differences in
23 growth performance in birds fed the PC or NC in experiments 2 or 3. In all three experiments,
24 birds fed the NC supplemented with neutral protease 1 had reduced ($P < 0.05$) feed intake
25 (**FI**) or body weight gain (**BWG**) and increased ($P < 0.05$) feed conversion ratio (**FCR**)
26 compared with birds fed the NC. Birds fed the NC diet supplemented with neutral protease 3,
27 7 (Experiment 1) or acid protease 4 (Experiment 3) had increased ($P < 0.05$) FCR and birds
28 fed neutral protease 6 (Experiment 2) had reduced ($P < 0.05$) BWG compared with birds fed
29 the NC. Apparent ileal amino acid digestibility was improved ($P < 0.05$) with protease
30 supplementation to the NC diets (Experiment 1 or 3), but this was dependent on the protease
31 and the amino acid. In conclusion, novel protease supplementation improved AID of amino
32 acids but this was not reflected in improvements in growth performance of turkey poultts or
33 broiler chicks.

34 **Keywords:** amino acids, broiler, turkey, protease, performance, apparent ileal digestibility

INTRODUCTION

35

36 The use of protease enzymes in industrial applications, such as detergents, textiles,
37 food processing and animal feed, is a major contributor to the \$5 billion market for industrial
38 enzymes (Juntunen et al., 2015). Proteases are classified into six groups: aspartate, cysteine,
39 glutamate, metallo, serine and threonine proteases, based on mechanistic features within each
40 group (Li et al., 2013). Over one-third of all known proteolytic enzymes are serine proteases
41 with an endoproteolytic catalytic activity typically dependent on a triad of aspartate, histidine
42 and serine residues (Di Cera, 2009). The endogenous proteases, trypsin and chymotrypsin,
43 belong to the largest family of serine proteases and cleave polypeptide chains at positively
44 charged arginine, lysine residues or large hydrophobic phenylalanine, tryptophan, tyrosine
45 residues, respectively (Di Cera, 2009). It has also been suggested that serine proteases are
46 allosteric enzymes and respond to the conditions of their environment differently, which may
47 influence biological activity and specificity (Di Cera, 2009), and impart differences in the
48 serine proteases selected for use in industrial applications.

49 Aspartic proteases are commonly called acid endopeptidases with aspartate residues at
50 their active site (Mandujan-Gonzalez et al., 2016). In the food industry, they are
51 predominantly used during the process of milk clotting to make cheese and to prevent
52 formation of wine haze (Schlander et al., 2017). Pepsin is a well known aspartic protease in
53 the A1 family with 282 other members (Dunn, 2002), and most aspartic proteases have broad
54 peptide bond specificity (Uniacke-Lowe and Fox, 2017). Swine pepsin and chicken pepsin
55 have similar molecular weights but contain different basic groups, and chicken pepsin has a
56 higher stability at alkaline solutions due to its smaller over-all negative charge (Bohak, 1969).
57 Esumi et al., (1980) reported the optimal pH values for quail and chicken pepsin were about
58 3.0, with quail pepsin having a higher relative activity at alkaline pH than chicken pepsin.
59 Therefore, within the same group of neutral serine or acidic aspartic endogenous or industrial

60 proteases, differences exist in their biological activity, substrate specificity, pH optima and
61 relative activity at a range of pH.

62 In animal feed, protease supplementation is of interest to improve protein and amino
63 acid digestibility, particularly in very young animals where the relative activity of
64 endogenous proteases may not be optimal (Lewis et al., 1955; Mahagna et al., 1995). In
65 addition, protease supplementation may improve ingredient quality by reducing ingredient
66 variability and mitigating negative effects of heat-stable trypsin-inhibitors or lectins
67 (Cowieson et al., 2016). Lewis et al. (1955) reported improvements in gain and efficiency of
68 piglets fed diets supplemented with pepsin, pancreatin, papain or a fungal protease from
69 *Aspergillus oryzae*. More recently, neutral serine or acid protease supplementation is gaining
70 in popularity in animal diets with beneficial (Angel et al., 2011; Cowieson and Roos, 2016)
71 or inconclusive (Freitas et al., 2011; Fru-Nji et al., 2011; Yuan et al., 2015; Yuan et al., 2017)
72 effects on growth performance, nutrient digestibility and endogenous enzyme secretion. The
73 objective of this set of trials was to evaluate the efficacy of eight neutral serine proteases and
74 six acid aspartic proteases when supplemented in low protein and amino acid diets fed to
75 turkey poults or broiler chicks for approximately 3 weeks from hatch. The response variables
76 included growth performance, apparent ileal digestibility of amino acids (AID), and
77 digestible amino acid intake in g/day.

78 MATERIALS AND METHODS

79 All animal procedures were approved by the Institutional Animal Care and Use
80 Committees of the University of Missouri (Experiments 1 and 2) and the National Institute of
81 Poultry Husbandry and approved by the Ethical Committee of Harper Adams University
82 (Experiment 3).

83 *Novel proteases*

84 The type of protease, source organism, working temperature optimum, pH optimum
85 and pH range of the evaluated proteases is listed in Table 1. There were 14 proteases
86 evaluated in 3 poultry trials, eight neutral proteases and six acid proteases. Eleven of the
87 proteases evaluated were novel proteases, and three proteases were obtained commercially.
88 In Experiments 1 and 2, all 14 proteases were evaluated. However, there was not enough
89 sample of acid protease 2 or 6 for inclusion in Experiment 3, and these proteases were
90 replaced with 1,500 or 3,000 FTU/kg of phytase (Quantum Blue, AB Vista, UK). Due to the
91 differences in pH optima, working temperature and substrate specificity, it was not possible
92 to standardize the dose supplemented in the experimental diets according to a specific unit/kg
93 or based on activity obtained from a universal assay. However, the amount of protease (mg)
94 in each sample was analyzed by determining the protease peak area obtained from HPLC
95 using a Superdex 75 10/300 GL column (GE Healthcare Bio-Sciences AB). Each protease
96 was diluted with wheat flour at different concentrations to allow for inclusion at an equivalent
97 enzyme protein concentration of 225 g in the final diet. This amount of enzyme protein was
98 obtained from the recommended dose of a commercially available serine protease and the
99 enzyme protein determined using the same assay as described above.

100 *Diets and experimental design*

101 Three separate experiments were conducted at two different Universities in the US or
102 UK using diet and husbandry conditions specific to each location. In each experiment, two
103 basal diets were formulated: a nutrient adequate positive control (**PC**), formulated to meet or
104 exceed the nutrient requirements for turkey poults (Experiment 1) or broiler chicks
105 (Experiments 2 and 3), and a negative control (**NC**) diet formulated to achieve 85%
106 (Experiments 1 and 2) or 80% (Experiments 3) of the requirement for protein and amino
107 acids. Phytase was included in all diets to provide 500 FTU/kg (Quantum Blue, AB Vista,
108 UK) and xylanase was included in all diets to provide 10,000 (Experiments 1 and 2) or

109 16,000 (Experiments 3) BXU/kg (Econase XT, AB Vista, UK). All diets were fed in mash
110 form, and birds were provided *ad libitum* access to feed and water throughout the duration of
111 the studies. Chromic oxide (Experiments 1 and 2) or titanium oxide (Experiment 3) was
112 added to the basal diets as an inert marker. The ingredient and nutrient composition of the
113 diets for Experiment 1, 2, and 3 are provided in Tables 2, 3 and 4, respectively.

114 ***Animals and husbandry***

115 ***Experiment one.*** Eight hundred, male, Hybrid Converter turkey poults were randomly
116 allocated to one of 16 experimental diets from one to 18 days post-hatch. Birds were housed
117 in Petersime battery brooder cages, with five birds/pen and 10 replicate pens/treatment. The
118 room temperature and humidity were thermostatically controlled and temperature maintained
119 at 29°C from d 1 to 7, 27°C from d 8 to 14, and 25°C from d 15 to 18. Light was provided to
120 the birds 24 hours/day for the duration of the study. Feed and water were provided in
121 troughs.

122 ***Experiment two.*** Eight hundred, male, Ross 308 broiler chicks were randomly
123 allocated to one of 16 experimental diets from one to 17-days post-hatch. Similar to
124 experiment 1, birds were housed in Petersime battery brooder cages with five birds/pen and
125 10 replicate pens/treatment. The room temperature and humidity were thermostatically
126 controlled and temperature maintained at 29°C from d 1 to 7, 27°C from d 8 to 14, and 25°C
127 from d 15 to 18. Light was provided to the birds 24 hours/day for the duration of the study.
128 Feed and water were provided in troughs.

129 ***Experiment three.*** Eight hundred, male, Ross 308 broiler chicks, vaccinated for
130 Marek's and Infectious Bursal disease at the hatchery, were randomly allocated to one of 16
131 experimental diets from one to 18-days post-hatch. Chicks were housed in metal battery
132 brooders on raised wire floors with five birds/pen and 10 replicate pens/treatment. The room
133 temperature was maintained with negative pressure ventilation, and gradually decreased from

134 32 to 22°C by the conclusion of the study. A standard lighting regime was used at 23:1 hours
135 light:dark from day old to 18:6 hours light:dark at day seven and maintained until the
136 conclusion of the trial.

137 *Sample collection, calculations and statistical analysis*

138 In all three experiments, feed and birds were weighed at the start and conclusion of
139 the trial to determine feed intake (**FI**) and BW gain (**BWG**) and calculate feed conversion
140 ratio (**FCR**). Birds were monitored daily and any culls or mortality was recorded to adjust
141 feed intake for bird days. At the conclusion of the studies, all birds were euthanized and ileal
142 digesta collected from the lower half of the ileum, defined as the section of intestine between
143 the Meckel's Diverticulum and the ileo-cecal junction. Digesta was pooled/pen,
144 homogenized, frozen and then dried for determination of amino acid and inert marker
145 concentration.

146 In Experiments 1 and 2, excreta were collected on the last 3 days of the trial,
147 pooled/pen, homogenized, frozen and dried prior to determination of starch and inert marker.
148 Diets, ileal digesta and excreta in Experiments 1 and 2 were analyzed for chromium (method
149 990.08) and diets and ileal digesta were analyzed for amino acids (method 982.30) according
150 to AOAC (2006). Starch was analyzed in the excreta samples using an enzymatic assay kit
151 (Sigma, St Louis, MO). Diets and ileal digesta in Experiment 3 were analyzed for titanium
152 according to methods of Short et al. (1996) and amino acids and starch according to
153 previously mentioned methods. To calculate digestible amino acid intake in g/day, the
154 following equation was used:

$$155 \quad \text{Digestible amino acid intake (g/d)} = (\text{analysed dietary amino acid (\%)} \times \\ 156 \quad \text{\% AID amino acid}) \times \text{daily intake (g)}$$

157 Data were analyzed as a two-way ANOVA using the fit model platform in JMP v.
158 13.0 (SAS, Cary, NC). Outliers were determined as three times the root mean square error

159 plus or minus the mean of response. Plotting the growth performance and AID data using a
160 normal quantile plot indicated the means were normally distributed. The model included
161 treatment and replicate pen. When treatment was significant, means were separated using
162 Dunnett's test for multiple comparisons. This test was used to compare each treatment to the
163 NC. Percent mortality was analyzed as nonparametric using a one-way ANOVA using the fit
164 Y by X platform in JMP v 13.0 (SAS, Cary, NC). If treatment was significant, differences
165 were established using the Steel's test for multiple comparisons (the non-parametric version
166 of the Dunnett's test). Significance was defined at $P < 0.05$, with trends discussed at $P <$
167 0.10 .

168 **RESULTS**

169 *Experiment one*

170 Nutrients, phytase and xylanase recoveries in the experimental diets were similar to
171 formulated values (Table 2). Overall mortality was 2.76% and not influenced by diet (Table
172 5). Poult fed the NC diet tended to gain less than those fed the PC ($P < 0.10$). Poult fed
173 neutral protease 1 ($P < 0.05$) ate less and gained less than poult fed the NC. Feed conversion
174 ratio tended to be higher in poult fed neutral protease 3 ($P < 0.10$) or neutral protease 7 ($P <$
175 0.10) when compared with those fed the NC. There were no other significant effects of diet
176 on growth performance in Experiment 1.

177 Apparent ileal digestibility of glutamate ($P < 0.10$), methionine ($P < 0.05$), isoleucine
178 ($P < 0.05$), leucine ($P < 0.10$), phenylalanine ($P < 0.10$) and arginine ($P < 0.10$) were greater
179 in poult fed the PC when compared with poult fed the NC (Table 6). Apparent ileal amino
180 acid digestibility of all measured amino acids was improved in poult fed neutral proteases 3
181 ($P < 0.05$) or 4 ($P < 0.10$), or acid proteases 2 ($P < 0.05$), 4 ($P < 0.05$) or 6 ($P < 0.10$) when
182 compared with poult fed the NC. Excluding tryptophan, AID of all other measured amino
183 acids was improved in poult fed neutral proteases 1 ($P < 0.05$), 2 ($P < 0.10$) or 6 ($P < 0.05$),

184 or acid protease 1 ($P < 0.05$) when compared with poult fed the NC. Poult fed neutral
185 protease 5 ($P < 0.10$) had improved AID of all amino acids measured except proline,
186 methionine, or tryptophan, when compared with poult fed the NC. Finally, poult fed
187 neutral protease 7 had improved glycine ($P < 0.10$), cysteine ($P < 0.10$), valine ($P < 0.05$),
188 isoleucine ($P < 0.10$), histidine ($P < 0.10$) and arginine ($P < 0.05$) digestibility, poult fed
189 neutral protease 8 had improved cysteine ($P < 0.05$), lysine ($P < 0.10$), and methionine ($P <$
190 0.05) digestibility, poult fed acid protease 3 had improved serine ($P < 0.05$) and histidine (P
191 < 0.05) digestibility, and poult fed acid protease 5 had improved histidine ($P < 0.10$)
192 digestibility when compared with poult fed the NC. Excreta starch retention was greater in
193 poult fed neutral protease 1 ($P < 0.05$) compared with poult fed the NC. There were no
194 other effects of treatment on excreta starch retention.

195 Poult fed the PC ($P < 0.05$) diet had a higher digestible amino acid intake for all
196 amino acids when compared with poult fed the NC diet (Table 7). Poult fed neutral
197 proteases 3 ($P < 0.05$) or 6 ($P < 0.05$) or acid proteases 1 ($P < 0.05$), 2 ($P < 0.05$), 3 ($P <$
198 0.05), 5 ($P < 0.05$) or 6 ($P < 0.05$) had a higher digestible amino acid intake for all amino
199 acids, except tryptophan, when compared with poult fed the NC. Poult fed neutral
200 proteases 2 ($P < 0.10$), 7 ($P < 0.10$) or 8 ($P < 0.05$) had increased digestible amino acid
201 intake, except for proline and/or tryptophan, when compared with poult fed the NC. Poult
202 fed neutral protease 5 ($P < 0.05$) or acid protease 4 ($P < 0.10$) had a higher digestible amino
203 acid intake, except for methionine or tryptophan, compared with poult fed the NC. Finally,
204 poult fed neutral protease 1 ($P < 0.05$) had reduced digestible amino acid intake of all amino
205 acids, except cysteine, when compared with poult fed the NC. There was no effect of
206 neutral protease 4 on digestible amino acid intake.

207 ***Experiment two***

208 Phytase and xylanase recoveries in the experimental diets were in agreement with
209 formulated values (Table 3). When analyzed in the diets, the protein and amino acid
210 concentration in the NC diet was only reduced by 6% compared with the PC (Table 3). This
211 was less than the expected 15% reduction from the PC, and may explain the non-significant
212 difference in FI, BWG or FCR of chicks fed the PC when compared with those fed the NC
213 (Table 8). Chicks fed neutral proteases 1, 4 or 6 ate significantly less and chicks fed neutral
214 proteases 1 or 6 gained significantly less than chicks fed the NC. There was no effect of diet
215 on FCR. Overall mortality was 4.0% and not influenced by diet.

216 There were very few effects of diet on AID of amino acids or excreta starch retention
217 (Table 9). Apparent ileal serine ($P < 0.10$) and lysine ($P < 0.05$) digestibility were improved
218 in chicks fed neutral protease 5 or acid protease 6, and methionine ($P < 0.05$) digestibility
219 was improved in chicks fed acid protease 5 or the PC when compared with chicks fed the NC.
220 There were no other effects of neutral protease supplementation on AID of amino acids.
221 Chicks fed acid proteases 1 ($P < 0.10$) or 3 ($P < 0.10$) had a lower AID of cysteine or
222 isoleucine compared with chicks fed the NC, and phenylalanine digestibility was reduced in
223 chicks fed acid protease 1 ($P < 0.05$) compared with chicks fed the NC. The AID of
224 tryptophan was reduced in chicks fed the PC ($P < 0.05$) compared with chicks fed the NC.
225 There were no other effects of acid protease supplementation on the AID of amino acids.
226 Apparent excreta starch retention was increased in chicks fed neutral proteases 1 ($P < 0.05$) or
227 3 ($P < 0.10$) compared with chicks fed the NC. There were no other effects of diet on starch
228 retention.

229 Similar to the AID data, chicks fed the PC had an increase in digestible methionine (P
230 < 0.05) intake and a decrease in digestible tryptophan ($P < 0.05$) intake when compared with
231 chicks fed the NC (Table 10). Chicks fed neutral protease 1 ($P < 0.05$) had a lower digestible
232 amino acid intake for all amino acids measured compared with chicks fed the NC. The

233 digestible amino acid intake of all measured amino acids, except lysine or lysine and
234 tryptophan, was lower in chicks fed neutral protease 6 ($P < 0.05$) or acid protease 3 ($P <$
235 0.05), respectively, compared with chicks fed the NC. Chicks fed neutral protease 4 ($P <$
236 0.10) had a reduced digestible intake of glutamate, proline, glycine, alanine, cysteine,
237 tyrosine, isoleucine, leucine, phenylalanine, histadine and tryptophan, and chicks fed neutral
238 protease 7 ($P < 0.10$) had reduced digestible amino acid intake of proline, tyrosine,
239 methionine, leucine, phenylalanine, and lysine compared with chicks fed the NC. There were
240 no other effects of neutral protease supplementation on digestible amino acid intake, except
241 for chicks fed neutral protease 5 ($P < 0.10$) having a higher digestible lysine and lower
242 digestible tryptophan intake compared with chicks fed the NC. Chicks fed acid protease 1 (P
243 < 0.10) had a lower digestible intake of most measured amino acids, except proline, lysine, or
244 tryptophan, compared with chicks fed the NC. Supplementation of the diets with acid
245 protease 4 ($P < 0.10$) reduced digestible methionine, phenylalanine, or tryptophan intake,
246 while acid protease 5 ($P < 0.10$) reduced digestible tyrosine or tryptophan intake and
247 increased digestible methionine intake compared with chicks fed the NC. There were no
248 other effects of acid protease supplementation on digestible amino acid intake.

249 *Experiment three*

250 Nutrient, phytase and xylanase recoveries in the experimental diets were as expected
251 and similar to formulated values (Table 4). Overall mortality was 2.4% and not influenced
252 by diet (Table 11). There were no differences in growth performance in chicks fed the PC
253 when compared with chicks fed the NC. Birds fed neutral protease 1 ($P < 0.01$) gained less
254 than birds fed the NC. Feed conversion ratio was higher in birds fed neutral protease 1 ($P <$
255 0.05) or acid protease 4 ($P < 0.10$) compared with birds fed the NC. There were no other
256 effects of diet on growth performance.

257 There was no difference in AID of amino acids or starch in birds fed the PC compared
258 with birds fed the NC (Table 12). There was no effect of acid protease or phytase
259 supplementation on the AID of any amino acids measured, or the AID of starch. The AID of
260 aspartate, serine, glutamate, glycine, alanine, threonine, valine, isoleucine, leucine,
261 phenylalanine, histidine and arginine were reduced in chicks fed neutral protease 8 ($P < 0.10$)
262 compared with chicks fed the NC. Apparent ileal serine or threonine digestibility were
263 reduced in chicks fed neutral proteases 3 ($P < 0.05$) or 7 ($P < 0.10$) when compared with
264 chicks fed the NC. Apparent ileal arginine or glycine digestibility were reduced ($P < 0.10$) in
265 chicks fed neutral proteases 4 ($P < 0.05$) or 7 ($P < 0.10$), respectively, when compared with
266 chicks fed the NC. Finally, the AID of serine, isoleucine, leucine, lysine, or histidine was
267 increased in chicks fed neutral protease 1 ($P < 0.10$) when compared with chicks fed the NC.
268 There were no other effects of diet on the AID of amino acids or starch.

269 Chicks fed the PC ($P < 0.05$) diet had a greater digestible amino acid intake compared
270 with chicks fed the NC. Contradictory to the AID of amino acids, chicks fed neutral protease
271 1 ($P < 0.10$) had a lower digestible intake of most measured amino acids, except serine,
272 glycine, tyrosine, methionine, isoleucine, or leucine, when compared with chicks fed the NC.
273 Chicks fed neutral protease 3 ($P < 0.10$) or acid protease 3 ($P < 0.05$) had a lower digestible
274 intake of all amino acids, except glutamate, tyrosine and methionine or proline and
275 methionine, respectively, when compared with chicks fed the NC. Supplementation of the
276 NC diet with acid protease 1 ($P < 0.10$) lowered the digestible intake of aspartate, proline,
277 glycine, alanine, cysteine, threonine, valine, phenylalanine, lysine, histidine and arginine
278 compared with chicks fed the NC. Finally, chicks fed the NC diet supplemented with neutral
279 proteases 2 ($P < 0.10$) or 4 ($P < 0.10$) or acid protease 5 ($P < 0.05$) had a lower digestible
280 intake of proline, glycine, cysteine, phenylalanine, histidine, and arginine or glycine, alanine,
281 cysteine, lysine, and arginine or aspartate, serine and cysteine, respectively, compared with

282 chicks fed the NC. Digestible intake of methionine was greater in chicks fed neutral
283 proteases 6 ($P < 0.05$), 7 ($P < 0.10$) or 8 ($P < 0.10$), acid protease 4 ($P < 0.05$) or phytase at
284 3000 FTU/kg ($P < 0.05$) compared with chicks fed the NC.

285 **DISCUSSION**

286 Each protease evaluated had specific pH and temperature optima and substrate
287 specificity. Protease recoveries in the experimental diets were not performed due to lack of
288 an acceptable in-feed assay that is universal, optimal or specific to each protease. All diets
289 were fed in mash form, and no denaturation of enzymatic activity would be expected in the
290 diets due to processing of the feed. Due to the number of experimental cages and the large
291 sample of proteases for testing, it was not possible to evaluate an optimum dose or dose
292 response in the current set of trials. However, three of the novel proteases (neutral proteases
293 1 and 5 and acid protease 5) were used in two subsequent trials to evaluate a dose response on
294 growth performance and apparent ileal amino acid digestibility in broilers (C. Walk et al.,
295 unpublished data). These trials evaluated doses that were lower, similar, and higher than the
296 dose evaluated in the current experiments and indicated the optimal dose of each protease
297 was similar or below (neutral protease 1) that of the dose employed in this set of trials (C.
298 Walk et al., unpublished data). Therefore, some of the detrimental effects reported from
299 supplementation the NC diet with neutral protease 1 may be associated with the dose selected
300 in the current trials.

301 Previous authors have reported significant improvements in growth performance
302 (Angel et al., 2011; Cowieson et al., 2016; Xu et al., 2017) or apparent ileal amino acid
303 digestibility of ingredients (Adebiyi and Olukosi, 2015; Stefanello et al., 2016) or diets
304 (Angel et al., 2011; Cowieson and Roos, 2014) supplemented with exogenous protease and
305 fed to broilers or turkeys. Others have reported no effect of supplemental protease on
306 performance, with a significant increase in apparent amino acid, protein or energy

307 digestibility (Freitas et al., 2011) or a reduction in performance and endogenous enzyme
308 activity as protease supplementation increased in the diet (Yuan et al., 2015; 2017).
309 Inconsistency in the effect of exogenous proteases on growth performance or amino acid
310 digestibility of poultry has been attributed to the inherent digestibility of amino acids in the
311 diets (Cowieson and Roos, 2016). In addition, variability in the source and quality of
312 soybean meal in the diet (Garcia-Rebollar et al., 2016) and protease enzymes that are not
313 clearly defined (Freitas et al., 2011) or supplemented in combination with other enzymes may
314 also contribute to the inconsistent reports surrounding protease supplementation in poultry
315 diets. The objective of these experiments was to evaluate the efficacy of novel serine or
316 aspartic proteases and three commercially available proteases on growth performance and
317 AID of amino acids in turkeys or broilers.

318 ***Performance***

319 To assess the potential efficacy of the novel proteases it was important that the NC
320 diet, the diet to which the test proteases were supplemented, was deficient in protein and
321 amino acids, to allow for noticeable improvements in growth performance, AID or digestible
322 amino acid intake. The reduction in amino acids and protein by 10 to 15% was modelled
323 after Angel et al. (2011), who reported a significant reduction in growth of chicks fed a low
324 protein and amino acid diet. In that experiment, protease supplementation improved growth
325 performance (Angel et al., 2011). In the current set of experiments, turkey poult fed the NC
326 diet tended to gain less than poult fed the PC (Experiment 1) but there was no effect of the
327 NC diet on growth performance of broiler chicks (Experiments 2 or 3). These results could
328 be expected in Experiment 2 with only a 6% reduction in the analyzed protein content
329 between the PC and NC diet. However, there was a 19% difference in protein and 28%
330 difference in total lysine between the PC and NC diets in Experiment 3, which would have
331 been expected to influence growth.

332 Regardless of the lack of an effect on growth performance between the PC and NC
333 diets in two of the three experiments, a few consistent responses were observed for the
334 different proteases evaluated. For example, in all three experiments birds fed neutral protease
335 1 ate less, gained less or were less efficient than birds fed the NC. Neutral protease 1 is an
336 extracellular subtilisin-like serine protease with commercial application in detergent
337 formulations (Juntunen et al., 2015). Juntunen et al. (2015) summarised the source organism
338 as belonging to a species of fungi frequently described as phytopathogenic. However, the
339 protease was expressed in *Trichoderma*, a commonly used organism for enzyme expression,
340 and therefore the source organism would have no effect on the actual protease that was fed in
341 the diet. More likely the significant reduction in growth performance of birds fed neutral
342 protease 1 were the result of an excess dose of the protease in the diet. Previous authors have
343 reported significant reductions in BWG as protease dose in the diet increased (Yuan et al.,
344 2017).

345 In experiment 1, FCR tended to be higher in poults fed neutral proteases 3 or 7 when
346 compared with poults fed the NC, and in experiment 3 acid protease 4 tended to increase
347 FCR compared with chicks fed the NC. Neutral protease 3 is classified as a proline-specific
348 endoprotease that can be used in the degradation of wheat gluten (Van Der Laan et al., 2017).
349 Neutral protease 7 is a commercially available serine-protease which has been previously
350 reported to improve feed efficiency of birds fed low protein and amino acid diets (Freitas et
351 al., 2011). Acid protease 4 is described as a pepsin-like protease, and previous authors have
352 reported significant improvements in rates of gain or feed efficiency with the
353 supplementation pepsin into piglet diets (Lewis et al., 1955; Baker, 1959). However, in a set
354 of subsequent experiments, Baker (1959) reported pepsin supplementation greater than 0.25%
355 or in diets containing dried skim milk reduced gain with no beneficial effects reported on
356 feed efficiency. Further, the author ran a series of experiments to determine factors that

357 influence pepsin efficacy and reported a 10% reduction in gain and 7% loss in feed efficiency
358 when pepsin was supplemented to low protein (15% CP) diets. Similarly, in the current trial
359 the differences in FCR are associated with a numeric increase in FI with less of an effect on
360 gain, and therefore the birds were able to eat through the protein deficiency but not utilize the
361 nutrients at an equivalent rate of gain.

362 ***Apparent ileal digestibility and digestible amino acid intake***

363 Baker (1959) reported 1.2 to 3.6% improvements in apparent protein digestibility in
364 piglets fed low protein diets supplemented with pepsin. Unfortunately, these improvements
365 in digestibility were not manifested as improvements in gain or efficiency in the low CP diet,
366 and the authors speculated this was related to feed passage rate, which will be discussed in
367 more detail below. The results in piglets presented by Baker (1959) and Freitas et al. (2011)
368 are in agreement with the results of the current set of trials, in which the improvement in AID
369 of amino acids (Experiment 1 or 3) was not associated with similar improvements in growth
370 performance. Previous authors have predicted protease supplementation will improve AID of
371 amino acids between 1.3 and 5.5%, with greater improvements noted when the control diet
372 amino acid digestibility is low (Cowieson and Roos, 2014). In the current set of trials, the
373 largest and most significant effect of protease supplementation was noted in experiment 1 in
374 which the average AID of amino acids in the NC diet was 80%, whereas in experiment 2 or 3
375 the average AID of amino acids in the NC diet was 86 and 88%, respectively. Therefore, as
376 previously reported, the AID of amino acids in the control diet will influence the magnitude
377 of the effect of protease supplementation on the digestibility of the diet. This may have
378 contributed to the lack of a significant effect of protease supplementation on AID of amino
379 acids in Experiment 2 or 3 but does not reflect the lack of an effect of protease on growth
380 performance, even with improvements in AID.

381 To try and understand the lack of an effect on growth performance with
382 improvements in the AID of amino acids, digestible amino acid intake in g/day was
383 calculated from the analysed amino acid in the diet, daily intake and the AID of the amino
384 acids. Protein deposition rate (or growth) should increase with increasing amino acid intake
385 and variations in intake can influence AID, with birds balancing their intake to fulfil
386 nutritional requirements (Cruz et al., 2005). However, the ability of the bird to adjust intake
387 based on nutrient requirements depends on the quality of the ingredients (Cruz et al., 2005) or
388 the digestible amino acids provide by the diet. Previous authors have reported pepsin
389 supplementation may exert beneficial effects on growth performance of piglets, partially
390 through a change in the rate of food passage with significant and positive correlations
391 between the rate of food passage and AID of protein (Baker, 1959). While the rate of food
392 passage was not measured in the current set of experiments, the digestible amino acid intake
393 in g/d may provide a better explanation for the lack of correlation between performance and
394 amino acid digestibility in the current experiments. For example, birds fed the PC diet had
395 significantly higher digestible amino acid intake (Experiments 1 and 3) compared with birds
396 fed the NC, while neutral protease 1 significantly improved AID of amino acids (Experiments
397 1 and 3) but was associated with a proportionately larger reduction in intake, resulting in
398 reduced digestible amino acid intake of all amino acids, hence the reduction in growth and
399 feed efficiency. Birds fed neutral protease 8 had significantly reduced AID of amino acids
400 (Experiment 3) in the absence of an effect on growth performance, possibly due to an
401 increase in digestible methionine intake, which was likely the most limiting amino acid in the
402 diet.

403 The digestible intake of all amino acids was increased in poult fed both neutral and
404 acid proteases, with the exception of neutral proteases 1 (as described earlier) or 4
405 (Experiment 1). However, in experiments 2 or 3, where there were less effects of protease

406 supplementation on AID of amino acids, protease supplementation had no effect or decreased
407 the digestible intake of amino acids compared with birds fed the NC, which may have
408 resulted in the lack of an effect of protease in these diets. This is contradictory to previously
409 published reports in similar diets (Angel et al., 2011), and may be indicative of imbalances in
410 the digestible amino acids available in the diet or alterations in endogenous protein digestion
411 due to an exogenous protease effect on endogenous proteolytic activity (Yuan et al. 2015;
412 2017).

413 *Conclusions*

414 In conclusion, the current set of trials evaluating the supplementation of 8 serine
415 proteases and 6 acid proteases failed to elicit beneficial effects of protease supplementation
416 on poultry growth performance, even in the presence of improvements in AID of amino
417 acids. This was associated with reductions in digestible amino acid intake. Further work to
418 evaluate a dose response of the novel proteases in the diets of poultry is ongoing.

419 **REFERENCES**

420 Adebiyi, A. O. and O. A. Olukosi. 2015. Apparent and standardised ileal amino acid
421 digestibility of wheat distillers dried grains with solubles with or without exogenous protease
422 in broilers and turkeys. Br. Poult. Sci. 56: <http://dx.doi.org/10.1080/00071668.2015.1011606>

423 Angel, C. R., W. Saylor, S. L. Vieira, and N. Ward. 2011. Effects of a
424 monocomponent protease on performance and protein utilization in 7- to 22-day-old broiler
425 chickens. Poult. Sci. 90:2281-2286.

426 AOAC. 2006. Official Methods of Analysis of AOAC International 18th ed.,
427 Arlington, VA.

428 Baker, R. O. 1959. Proteolytic enzymes in baby pig nutrition. Ph.D. Thesis. Iowa
429 State University, Ames.

430 Bohak, Z. 1969. Purification and characterization of chicken pepsinogen and chicken
431 pepsin. *J. Biol. Chem.* 244:4638-4648.

432 Cowieson, A. J., H. Lu, K. M. Ajuwon, I. Knap, and O. Adeola. 2016. Interactive
433 effects of dietary protein source and exogenous protease on growth performance, immune
434 competence and jejunal health of broiler chickens. *Anim. Prod. Sci.* 57:252-261.

435 Cowieson, A. J. and F. F. Roos. 2016. Toward optimal value creation through the
436 application of exogenous mono-component proteases in the diets of non-ruminants. *Anim.*
437 *Feed Sci. Technol.* 221:331-340.

438 Cruz, V. C., Pezzato, A. C., Pinheiro, D. F., Goncalves, J. C., and Sartori, J. R. 2005.
439 Effect of free-choice feeding on the performance and ileal digestibility of nutrients in
440 broilers. *Braz. J. Poult. Sci.* 7:143-150.

441 Di Cera, E. 2009. Serine proteases. *IUBMB Life.* 61:510-515.

442 Dunn, B. M. 2002. Structure and mechanism of the pepsin-like family of aspartic
443 peptidases. *Chem. Rev.* 102:4431-4458.

444 Esumi, H., S. Yasugi, T. Mizuno, and H. Fujiki. 1980. Purification and
445 characterization of a pepsinogen and its pepsin from the proventriculus of the Japanese quail.
446 *Biochimica. Biophysica. Acta.* 611:363-370.

447 Freitas, D. M., S. L. Vieira, C. R. Angel, A. Favero, and A. Maiorka. 2011.
448 Performance and nutrient utilization of broilers fed diets supplemented with a novel mono-
449 component protease. *J. Appl. Poult. Res.* 20:322-334.

450 Fru-Nji, F., A. M. Kluentner, M. Fischer, and K. Pontoppidan. 2011. A feed serine
451 protease improves broiler performance and increases protein and energy digestibility. *J.*
452 *Poult. Sci.* 48:239-246.

453 Garcia-Rebollar, P., L. Camara, R. P. Lazaro, C. Dapoza, R. Perez-Maldonado, and G.
454 G. Mateos. 2016. Influence of the origin of the beans on the chemical composition and
455 nutritive value of commercial soybean meals. *Anim. Feed Sci. Technol.* 221:245-261.

456 Ireta, J. M. and L. S. Gilchrist. 1994. Fusarium Head Scab of Wheat (*Fusarium*
457 *graminearum* Schwabe). Wheat Special Report No. 21b. Mexico, D. F.: CIMMYT.
458 ISBN: 968-6923-21-7.

459 Lewis, C. J., D. V. Catron, C. H. Liu, V. C. Speer, and G. C. Ashton. 1955. Enzyme
460 supplementation of baby pig diets. *Agric. Food Chem.* 3:1047-1050.

461 Li, Q., L. Yi, P. Marek, and B. L. Iverson. 2013. Commercial proteases: Present and
462 future. *FEBS Letters.* 587:1155-1163.

463 Mahagna, M. I. Nir, M. Larbier, and Z. Nitsan. 1995. Effect of age and exogenous
464 amylase and protease on development of the digestive tract, pancreatic enzyme activities and
465 digestibility of nutrients in young meat-type chicks. *Reprod. Nutr. Dev.* 35:201-212.

466 Mandujano-Gonzalez, V., L. Villa-Tanaca, M. A. Anducho-Reyes, Y. Mercado-
467 Flores. 2016. Secreted fungal aspartic proteases: A review. *Rev. Iberoam. Micol.* 33:76-82.

468 Schlander, M. U. Distler, S. Tenzer, E. Thines, and H. Claus. 2017. Purification and
469 properties of yeast proteases secreted by *Wickerhamomyces anomalus* 227 and *Metschnikovia*
470 *pulcherrima* 446 during growth in white grape juice. *Fermentation.* 3:
471 doi:10.3390/fermentation3010002

472 Short, F. J., P. Gordon, J. Wiseman, and K. N. Boorman. 1996. Determination of
473 titanium dioxide added as an inert marker in chicken digestibility studies. *Anim. Feed Sci.*
474 *Tech.* 59:215-221.

475 Stefanello, C., S. L. Vieira, H. V. Rios, C. T. Simoes, and J. O. B. Sorbara. 2016.
476 Energy and nutrient utilisation of broilers fed soybean meal from two different Brazilian
477 production areas with an exogenous protease. *Anim. Feed Sci. Technol.* 221:267-273.

478 Uniacke-Lowe, T. and P. F. Fox. 2017. Chapter 4 - Chymosin, Pepsins and Other
479 Aspartyl Proteinases: Structure, Functions, Catalytic Mechanism and Milk-Clotting
480 Properties in Cheese. Chemistry, Physics and Microbiology pg. 69-113. 4th ed. P. L. H.
481 McSweeney, P. F. Fox, P. Cotter, and D. W. Everett. Academic Press, Cambridge
482 Massachusetts, US.

483 Van Der Laan, J. M., P. J. I. Van De Vondervoort, C. Christis, M. Spaans, A. De
484 Bruine-Paulus, J. H. M. Mutsaers. 2017. US Patent No. 20170081651 A1. Proline specific
485 endoprotease. DSM IP Assets B.V., Netherlands.

486 Xu, X., H. L. Wang, L. Pan, X. K. Ma, Q. Y. Tian, Y. T. Xu, S. F. Long, Z. H. Zhang,
487 and X. S. Piao. 2017. Effects of coated proteases on the performance, nutrient retention, gut
488 morphology and carcass traits of broilers fed corn or sorghum based diets supplemented with
489 soybean meal. Anim. Feed. Sci. Technol. 223:119-127.

490 Yuan, L. S. Q. Wang, Z. X. Wang, H. Zhu, and K. Huang. 2015. Effects of exogenous
491 protease supplementation on endogenous trypsin activity and gene expression in broilers.
492 Genet. Molec. Res. 14:13633-13641.

493 Yuan, L., M. Wang, X. Zhang, and Z. Wang. 2017. Effects of protease and non-starch
494 polysaccharide enzyme on performance, digestive function, activity and gene expression of
495 endogenous enzymes of broilers. PLOS One.12: e0173941.

496 **Table 1.** Description of novel proteases evaluated

Category	#	pH (optimum)	Temperature ¹ , °C (optimum)	Source organism	Protease type	Experiment ²
Neutral						
	1	6 – 11 (10)	40 – 65 (60)	<i>Fusarium equiseti</i>	Subtilisin-like serine protease	1, 2, 3
	2	6 – 11 (10)	55 – 70 (70)	<i>Malbranchea cinnamomea</i>	Subtilisin-like serine protease	1, 2, 3
	3	6 – 8.5 (7)	40 – 50 (50)	<i>Myricoccum thermophilum</i>	Proline-specific endoprotease	1, 2, 3
	4	6 – 10 (9)	30 – 55 (50)	<i>Trichoderma reesei</i>	Subtilisin-like serine protease	1, 2, 3
	5	6 – 9 (8)	ND	<i>Trichoderma reesei</i>	Serine	1, 2, 3
	6	6 – 11 (9)	40 – 65 (60)	<i>Verticillium dahlia</i>	Subtilisin-like serine protease	1, 2, 3
	7	7 – 10	60 – 80 (70)	<i>Nocardiosis prasina</i>	Serine-specific protease	1, 2, 3
	8	7 – 10	60 – 80 (70)	<i>Bacillus licheniformis</i>	Serine	1, 2, 3
Acid						
	1	4 – 6.5 (6)	20 – 45	<i>Trichoderma reesei</i>	Subtilisin-like serine protease	1, 2, 3
	2	4 – 6 (5)	ND	<i>Trichoderma reesei</i>	Serine	1, 2
	3	3 – 8 (5 – 7)	< 50 (40)	<i>Trichoderma reesei</i>	Pepsin	1, 2, 3
	4	3 – 8 (5 – 6)	40 – 70 (60)	<i>Trichoderma reesei</i>	Pepsin	1, 2, 3
	5	2 – 5.5 (5)	(50)	<i>Trichoderma reesei</i>	Aspartyl	1, 2, 3
	6	ND ³ (acidic)	ND	<i>Streptomyces</i>		1, 2

497 ¹ Working temperature rather than thermostable temperature.498 ² There was not enough sample of acid protease 2 or 6 to test in all 3 experiments. For consistency, the protease numbers were maintained in
499 Experiment 3 without acid protease 2 or 6.500 ³ Not determined.

501 **Table 2.** Ingredient and nutrient composition of the starter
 502 diets, as-fed basis (Experiment 1)

Ingredient, %	Positive control	Negative control
Corn	43.59	55.59
Soybean meal	48.54	37.95
Corn oil	1.67	0.21
Salt	0.26	0.26
Sodium bicarbonate	0.17	0.17
DL methionine	0.33	0.25
Lysine HCl	0.30	0.32
Threonine	0.04	0.05
Limestone	1.25	1.27
Dicalcium phosphate	2.89	2.95
Phytase ¹	0.01	0.01
Vitamin premix ²	0.12	0.12
Trace mineral premix ³	0.13	0.13
Space (enzyme)	0.00	0.60
Chromic oxide	0.10	0.10
Xylanase ⁴	0.006	0.006
Nutrient composition, %		
Crude protein	27.50	23.38
ME, kcal/kg	2865	2865
Calcium	1.29	1.29
Phosphorus	1.03	1.00
Available phosphorus	0.65	0.65
Crude fat	3.99	2.89
Crude fiber	2.70	2.62
Methionine	0.74	0.61
Cysteine	0.43	0.38
TSAA	1.17	0.99
Lysine	1.80	1.53
Tryptophan	0.33	0.27
Threonine	1.10	0.94
Arginine	1.90	1.57
Phytate phosphorus	0.25	0.23
Sodium	0.17	0.17
Chloride	0.25	0.25
Analysed nutrients, %		
Crude protein	27.45	23.07
Total lysine	1.76	1.56
Total threonine	1.01	0.89
Total methionine	0.71	0.52
Phytase, FTU/kg	710	574
Xylanase, BXU/kg	17,600	13,413

503 ¹ Quantum Blue 5G (AB Vista) included to provide 500 FTU/kg.

504 ² Supplied per kilogram of diet: vitamin A, 7700 IU; vitamin D3, 2750 IU; vitamin E, IU

505 16.5; vitamin B₁₂, 11 ug; vitamin K, 0.83 mg; riboflavin, 6.6 mg; thiamin, 1.1 mg;
506 pantothenic acid, 6.6 mg; niacin, 27.5 mg; pyridoxine, 1.37 mg; folic acid, 0.69 mg; biotin,
507 33 mg; choline, 385 mg.
508 ³ Supplied per kilogram of diet: manganese (manganese sulfate), 100 mg; zinc (zinc oxide),
509 100 mg; iron (ferrous sulfate), 50 mg; cupper (copper sulfate), 11.25 mg; iodine (calcium
510 iodate), 1.5 mg; selenium (sodium selenite), 0.15 mg.
511 ⁴ Econase XT 25G (AB Vista) included to provide 9,600 BXU/kg.

512 **Table 3.** Ingredient and nutrient composition of the starter
 513 diets, as-fed basis (Experiment 2)

Ingredient, %	Positive control	Negative control
Corn	57.18	67.25
Soybean meal	36.85	27.95
Corn oil	1.40	0.17
Salt	0.29	0.29
Sodium bicarbonate	0.17	0.17
DL methionine	0.35	0.29
Lysine HCl	0.25	0.28
Threonine	0.09	0.08
Tryptophan	0.00	0.01
Limestone	1.14	1.16
Dicalcium phosphate	1.32	1.37
Phytase ¹	0.01	0.01
Vitamin premix ²	0.12	0.12
Trace mineral premix ³	0.13	0.13
Space (enzyme)	0.00	0.60
Chromic oxide	0.10	0.10
Xylanase ⁴	0.006	0.006
Nutrient composition, %		
Crude protein	23.00	19.55
ME, kcal/kg	3000	3000
Calcium	0.85	0.85
Phosphorus	0.66	0.64
Available phosphorus	0.35	0.35
Crude fat	4.12	3.19
Crude fiber	2.63	2.56
Methionine	0.71	0.60
Cysteine	0.37	0.33
TSAA	1.08	0.93
Lysine	1.44	1.22
Tryptophan	0.27	0.23
Threonine	0.97	0.82
Arginine	1.54	1.26
Phytate phosphorus	0.23	0.22
Sodium	0.18	0.18
Chloride	0.26	0.26
Analysed nutrients, %		
Crude protein	20.40	19.13
Total lysine	1.30	1.31
Total threonine	0.84	0.79
Total methionine	0.58	0.53
Phytase, FTU/kg	652	575
Xylanase, BXU/kg	14,700	13,893

514 ¹ Quantum Blue 5G (AB Vista) included to provide 500 FTU/kg.

515 ² Supplied per kilogram of diet: vitamin A, 7700 IU; vitamin D3, 2750 IU; vitamin E, IU
516 16.5; vitamin B₁₂, 11 ug; vitamin K, 0.83 mg; riboflavin, 6.6 mg; thiamin, 1.1 mg;
517 pantothenic acid, 6.6 mg; niacin, 27.5 mg; pyridoxine, 1.37 mg; folic acid, 0.69 mg; biotin,
518 33 mg; choline, 385 mg.

519 ³ Supplied per kilogram of diet: manganese (manganese sulfate), 100 mg; zinc (zinc oxide),
520 100 mg; iron (ferrous sulfate), 50 mg; copper (copper sulfate), 11.25 mg; iodine (calcium
521 iodate), 1.5 mg; selenium (sodium selenite), 0.15 mg.

522 ⁴ Econase XT 25G (AB Vista) included to provide 9,600 BXU/kg.

523 **Table 4.** Ingredient and nutrient composition of the starter
 524 diets, as-fed basis (Experiment 3)

Ingredient, %	Positive control	Negative control
Wheat	59.20	70.67
Soybean meal	32.36	20.74
Soy oil	4.10	3.43
Salt	0.25	0.25
Sodium bicarbonate	0.19	0.18
DL methionine	0.36	0.32
Lysine HCl	0.36	0.49
Threonine	0.13	0.18
Tryptophan	0.00	0.02
Limestone	1.21	1.27
Dicalcium phosphate	1.02	1.03
Phytase ¹	0.01	0.01
Vitamin mineral premix ²	0.50	0.50
Space (enzyme)	0.00	0.60
Titanium oxide	0.30	0.30
Xylanase ⁴	0.01	0.01
Nutrient composition, %		
Crude protein	22.50	18.25
ME, kcal/kg	3025	3025
Calcium	0.85	0.85
Phosphorus	0.63	0.59
Available phosphorus	0.35	0.35
Crude fat	5.39	4.83
Crude fiber	2.55	2.44
Methionine	0.68	0.58
Cysteine	0.38	0.32
TSAA	1.06	0.90
Lysine	1.43	1.22
Tryptophan	0.28	0.24
Threonine	0.93	0.79
Arginine	1.44	1.08
Phytate phosphorus	0.23	0.20
Sodium	0.18	0.18
Chloride	0.27	0.29
Analysed nutrients, %		
Crude protein	22.10	17.95
Total lysine	1.69	1.22
Total threonine	1.04	0.79
Total methionine	0.77	0.55
Phytase, FTU/kg	511	549
Xylanase, BXU/kg	17,600	19,600

525 ¹ Quantum Blue 5G (AB Vista) included to provide 500 FTU/kg.

526 ² Supplied the following per kilogram of diet: vitamin A, 5,484 IU; vitamin D₃, 2,643 ICU;
527 vitamin E, 11 IU; menadione sodium bisulfite, 4.38 mg; riboflavin, 5.49 mg; d-pantothenic
528 acid, 11 mg; niacin, 44.1 mg; choline chloride, 771 mg; vitamin B12, 13.2 µg; biotin, 55.2
529 µg; thiamine mononitrate, 2.2 mg; folic acid, 990 µg; pyridoxine hydrochloride, 3.3 mg; I,
530 1.11 mg; Mn, 66.06 mg; Cu, 4.44 mg; Fe, 44.1 mg; Zn, 44.1 mg; Se, 250 µg.
531 ³ Econase XT 25G (AB Vista) included to provide 16,000 BXU/kg.

532 **Table 5.** Growth performance of turkey poult fed reduced nutrient density diets
 533 and novel proteases from hatch to 18 days post-hatch (Experiment 1)

Diet	Protease	Feed intake, g	BW gain, g	FCR, g:g	Mortality, %
Negative control		708.7	494.3	1.434	12.5
	Neutral				
	1	512.5*	344.7*	1.499	5.0
	2	719.7	473.5	1.523	0.0
	3	749.6	480.8	1.559 [†]	0.0
	4	685.3	463.6	1.524	4.0
	5	730.3	507.5	1.439	0.0
	6	726.1	490.2	1.486	6.0
	7	747.9	483.7	1.551 [†]	0.0
	8	746.8	491.8	1.520	4.0
	Acid				
	1	745.9	487.9	1.529	0.0
	2	727.5	487.7	1.498	5.0
	3	748.5	486.0	1.502	0.0
	4	694.6	469.2	1.490	4.4
	5	730.8	496.3	1.475	2.2
	6	724.1	480.7	1.513	2.0
	Positive control	774.1	546.3 [†]	1.417	0.0
	SE	18.3	12.9	0.03	0.03
	Diet P-value	< 0.0001	< 0.0001	0.0316	0.12
	Replicate P-value	0.13	0.0003	0.0081	0.31

534 Means in the same column are significantly different from the negative control, *P
 535 < 0.05, [†]P < 0.10.

536 **Table 6.** Apparent ileal amino acid digestibility and apparent excreta starch retention of turkey poult fed reduced crude protein and amino acid
 537 diets and novel proteases from hatch to 18 days post-hatch (Experiment 1)

Nutrient	NC ¹	Neutral protease								Acid protease						PC ²	SE
		1	2	3	4	5	6	7	8	1	2	3	4	5	6		
NEAA ³																	
Asp	80.6	84.9*	85.0*	86.9*	85.0*	85.3*	87.1*	83.5	82.7	85.6*	85.8*	83.4	86.8*	83.1	86.1*	83.6	0.9 ^{5,6}
Ser	80.5	84.6*	84.2 [†]	86.4*	84.6*	84.9*	87.6*	83.4	81.9	85.1*	85.3*	85.1*	87.0*	82.3	85.5*	83.5	0.9 ^{5,6}
Glu	83.5	88.5*	87.0*	89.2*	87.5*	87.2*	89.6*	86.3	85.6	88.0*	88.9*	86.2	89.4*	85.5	88.7*	86.7 [†]	0.9 ^{5,6}
Pro	78.7	83.8*	82.6 [†]	85.8*	82.9 [†]	82.4	85.5*	81.1	80.5	83.4*	84.7*	81.0	85.6*	80.5	84.4*	81.9	1.1 ⁵
Gly	75.8	81.5*	81.7*	84.3*	81.4*	81.8*	83.9*	79.7 [†]	79.3	82.3*	82.7*	79.4	83.6*	78.6	82.5*	79.7	1.0 ^{5,6}
Ala	77.0	83.2*	81.4 [†]	85.2*	82.5*	81.5 [†]	84.2*	79.7	79.7	82.5*	83.6*	80.1	84.2*	78.0	82.8*	80.4	1.2 ⁵
Cys	63.9	73.2*	74.7*	77.0*	71.9*	72.7*	76.5*	69.9 [†]	70.5*	75.9*	74.4*	68.7	74.5*	69.0	74.6*	69.9	1.6 ⁵
Tyr	78.6	85.2*	83.7*	86.8*	84.3*	83.5*	86.3*	81.5	81.1	84.3*	85.6*	81.5	85.8*	80.7	84.6*	82.0	1.0 ⁵
EAA ⁴																	
Thr	75.9	80.8*	81.2*	83.3*	80.6*	81.2*	83.0*	79.2	78.2	81.1*	81.3*	79.4	82.9*	77.5	82.0*	79.3	1.0 ⁵
Val	76.6	84.6*	82.7*	84.7*	81.8*	81.4*	83.8*	81.0*	80.0	82.6*	82.6*	79.2	84.2*	79.2	83.0*	80.4	1.0 ⁵
Met	88.0	92.5*	91.6*	93.0*	91.3*	90.1	92.9*	90.0	91.1*	91.4*	92.2*	90.0	92.1*	89.5	92.1*	92.6*	0.7 ⁵
Iso	80.2	86.1*	84.9*	87.4*	84.9*	84.7*	86.4*	83.6 [†]	83.1	85.5*	86.3*	82.3	86.6*	82.2	86.0*	84.1*	0.9 ⁵
Leu	79.6	86.3*	83.7 [†]	86.9*	84.5*	83.5 [†]	86.3*	82.6	82.2	84.6*	86.2*	82.4	86.4*	81.2	85.6*	83.5 [†]	1.0 ⁵
Phe	81.7	87.6*	85.8*	88.2*	86.2*	85.8*	88.0*	84.8	84.5	86.4*	87.7*	84.3	87.8*	83.8	87.1*	85.3 [†]	0.9 ⁵
Lys	87.1	90.2*	90.4*	91.2*	90.3*	90.6*	90.9*	89.1	89.2 [†]	90.4*	90.4*	89.0	91.0*	88.9	90.8*	89.1	0.6 ⁵
His	84.1	89.2*	87.6*	89.8*	88.1*	87.5*	89.5*	86.9 [†]	86.4	88.2*	89.3*	86.3	89.6*	86.1	88.7*	86.7	0.8 ⁵
Arg	88.0	92.4*	91.0*	92.4*	91.1*	91.4*	93.1*	90.9*	90.2	91.6*	92.4*	90.7*	92.4*	90.3 [†]	92.3*	90.6 [†]	0.6 ^{5,6}
Trp	87.2	88.4	89.3	90.8*	89.6 [†]	89.2	88.7	88.3	88.8	89.3	90.4*	87.9	90.2*	87.4	89.6 [†]	89.3	0.7 ^{5,6}
Starch	85.0	94.1*	82.9	84.8	85.9	87.3	89.8	84.0	82.8	81.6	87.4	84.1	85.4	87.7	87.0	85.7	1.6 ^{5,6}

538 ¹ Reduced nutrient density negative control.

539 ² Nutrient adequate positive control.

540 ³ Non-essential amino acids.

541 ⁴ Essential amino acids.

542 ⁵ Significant effect of diet (P < 0.05). Means in the same column are significantly different from the negative control, *P < 0.05, [†]P < 0.10.

543 ⁶ Significant effect of block (P < 0.05).

544 **Table 7.** Apparent digestible amino acid intake (g/day) of turkey poults fed reduced crude protein and amino acid diets and novel proteases from
 545 hatch to 18 days post-hatch (Experiment 1)

Nutrient	NC ¹	Neutral protease								Acid protease						PC ²	SE
		1	2	3	4	5	6	7	8	1	2	3	4	5	6		
NEAA ³																	
Asp	0.58	0.46*	0.69*	0.71*	0.62	0.71*	0.74*	0.69*	0.70*	0.71*	0.71*	0.72*	0.69*	0.72*	0.71*	0.83*	0.02 ⁵
Ser	0.24	0.19*	0.28 [†]	0.29*	0.26	0.30*	0.32*	0.28*	0.28*	0.29*	0.29*	0.33*	0.28*	0.28*	0.28*	0.33*	0.01 ⁵
Glu	1.02	0.80*	1.17*	1.24*	1.09	1.21*	1.27*	1.20*	1.21*	1.23*	1.23*	1.25*	1.19*	1.24*	1.22*	1.40*	0.03 ⁵
Pro	0.32	0.24*	0.36	0.38*	0.33	0.37*	0.39*	0.36	0.36	0.38*	0.37*	0.37*	0.36 [†]	0.37*	0.37*	0.41*	0.01 ⁵
Gly	0.22	0.18*	0.26*	0.27*	0.24	0.27*	0.28*	0.26*	0.27*	0.27*	0.27*	0.27*	0.26*	0.27*	0.27*	0.31*	0.01 ⁵
Ala	0.27	0.21*	0.31 [†]	0.33*	0.29	0.32*	0.33*	0.31*	0.31*	0.33*	0.32*	0.32*	0.31*	0.31*	0.32*	0.36*	0.01 ⁵
Cys	0.06	0.06	0.08*	0.09*	0.07	0.08*	0.09*	0.08*	0.09*	0.09*	0.08*	0.08*	0.08*	0.09*	0.09*	0.09*	0.00 ⁵
Tyr	0.17	0.13*	0.20*	0.22*	0.19	0.20*	0.21*	0.19	0.20*	0.21*	0.21*	0.20*	0.20*	0.20*	0.20*	0.22*	0.01 ⁵
EAA ⁴																	
Thr	0.21	0.17*	0.25*	0.26*	0.23	0.26*	0.27*	0.25*	0.25*	0.25*	0.26*	0.26*	0.25*	0.25*	0.26*	0.29*	0.01 ⁵
Val	0.26	0.21*	0.31*	0.32*	0.28	0.31*	0.32*	0.31*	0.32*	0.32*	0.31*	0.30*	0.31*	0.32*	0.32*	0.36*	0.01 ⁵
Met	0.14	0.11*	0.17*	0.16*	0.14	0.15	0.18*	0.16 [†]	0.19*	0.17*	0.17*	0.17*	0.15	0.17*	0.18*	0.24*	0.00 ⁵
Iso	0.24	0.19*	0.29*	0.30*	0.26	0.29*	0.30*	0.29*	0.30*	0.29*	0.29*	0.28*	0.29*	0.30*	0.30*	0.34*	0.01 ⁵
Leu	0.49	0.39*	0.56*	0.60*	0.53	0.57*	0.60*	0.57*	0.57*	0.59*	0.59*	0.59*	0.57*	0.58*	0.58*	0.66*	0.02 ⁵
Phe	0.28	0.23*	0.33*	0.35*	0.30	0.34*	0.36*	0.34*	0.34*	0.34*	0.35*	0.35*	0.34*	0.35*	0.34*	0.40*	0.01 ⁵
Lys	0.42	0.32*	0.49*	0.48*	0.44	0.52*	0.49*	0.49*	0.52*	0.49*	0.49*	0.51*	0.48*	0.52*	0.50*	0.58*	0.01 ⁵
His	0.17	0.13*	0.19*	0.20*	0.18	0.19*	0.20*	0.20*	0.20*	0.20*	0.20*	0.20*	0.19*	0.20*	0.20*	0.23*	0.01 ⁵
Arg	0.40	0.31*	0.47*	0.48*	0.42	0.47*	0.50*	0.47*	0.48*	0.48*	0.49*	0.48*	0.46*	0.49*	0.48*	0.56*	0.01 ⁵
Trp	0.08	0.06*	0.09	0.09	0.08	0.09	0.08	0.08	0.09*	0.09	0.09	0.09 [†]	0.08	0.08	0.09	0.10*	0.00 ⁵

546 ¹ Reduced nutrient density negative control.

547 ² Nutrient adequate positive control.

548 ³ Non-essential amino acids.

549 ⁴ Essential amino acids.

550 ⁵ Significant effect of diet (P < 0.05). Means in the same column are significantly different from the negative control, *P < 0.05, [†]P < 0.10.

551 ⁶ Significant effect of block (P < 0.05).

552 **Table 8.** Growth performance of broiler chicks fed reduced nutrient density diets
 553 and novel proteases from hatch to approximately 17 days post-hatch (Experiment 2)

Diet	Protease	Feed intake, g	BW gain, g	FCR, g:g	Mortality, %
Negative control		724.8	525.4	1.388	0.0
	Neutral				
	1	461.0*	332.7*	1.388	6.0
	2	666.1	485.4	1.389	2.0
	3	684.3	493.9	1.357	4.0
	4	639.3*	466.7	1.373	6.0
	5	679.3	481.5	1.418	8.0
	6	602.1*	439.8*	1.377	10.0
	7	675.1	471.4	1.384	4.0
	8	718.2	523.6	1.366	2.0
	Acid				
	1	690.1	477.8	1.448	2.0
	2	689.9	490.8	1.398	6.0
	3	669.5	461.7	1.438	2.0
	4	666.2	463.6	1.432	6.0
	5	671.2	485.8	1.391	0.0
	6	668.4	481.8	1.402	4.0
	Positive control	710.8	540.9	1.332	2.0
	SE	18.3	16.2	0.03	0.03
	Diet P-value	< 0.0001	< 0.0001	0.52	0.75
	Block P-value	0.0271	0.0100	0.10	0.71

554 Means in the same column are significantly different from the negative control, *P <
 555 0.05, †P < 0.10.

556 **Table 9.** Apparent ileal amino acid digestibility and apparent excreta starch retention of broiler chicks fed reduced crude protein and amino acid
 557 diets and novel proteases from hatch to 17 days post-hatch (Experiment 2)

Nutrient	NC ¹	Neutral protease								Acid protease						PC ²	SE
		1	2	3	4	5	6	7	8	1	2	3	4	5	6		
NEAA ³																	
Asp	85.6	85.3	86.5	85.8	85.5	87.4	84.7	86.6	86.9	83.7	86.5	84.0	86.3	87.1	87.8	84.5	0.7 ⁵
Ser	86.9	87.9	88.7	88.6	87.5	89.5 [†]	86.7	89.0	88.9	86.1	88.4	87.4	88.1	88.7	89.4 [†]	88.4	0.7 ^{5,6}
Glu	92.1	91.7	92.1	92.6	92.2	92.6	91.6	92.0	92.4	91.4	93.2	91.6	92.2	92.6	93.5	91.9	0.5
Pro	84.3	86.1	85.2	84.8	84.2	85.2	82.7	83.4	85.6	83.1	84.3	81.9	83.7	86.1	85.1	84.8	0.7 ⁵
Gly	82.0	81.3	81.8	81.3	80.7	83.4	80.6	82.3	83.8	79.8	82.6	79.7	82.6	82.9	83.4	82.2	0.7 ⁵
Ala	82.6	83.6	82.8	82.4	81.9	83.8	81.5	83.0	84.4	80.9	82.8	80.4	82.9	83.5	84.7	83.9	0.8 ⁵
Cys	78.9	79.7	79.3	78.9	77.6	80.0	77.3	78.1	80.4	75.0 [†]	79.1	73.8 [*]	78.7	80.4	80.2	77.8	1.0 ⁵
Tyr	84.6	87.8	86.9	87.2	86.2	84.5	83.1	84.1	84.6	81.6	86.7	83.2	85.2	84.4	85.6	85.9	0.9 ⁵
EAA ⁴																	
Thr	82.1	81.0	81.6	81.3	81.0	84.0	81.1	83.7	84.4	80.6	82.8	80.2	83.3	83.3	84.3	83.8	0.7 ⁵
Val	81.7	81.9	82.1	81.5	81.4	83.5	80.9	82.2	83.6	79.9	82.5	80.0	82.9	83.6	83.7	82.0	0.7 ⁵
Met	93.6	94.7	94.5	94.5	93.8	94.3	93.2	93.6	94.8	93.2	94.0	93.2	93.2	95.1 [*]	94.8	95.0 [*]	0.4 ⁵
Iso	85.3	84.5	84.3	84.0	84.1	86.7	84.2	85.7	86.4	83.1 [†]	86.0	83.1 [†]	85.9	86.4	86.9	86.0	0.6 ⁵
Leu	86.1	86.6	85.5	85.6	85.2	86.6	84.8	85.7	87.1	83.7	86.5	85.0	86.0	86.8	87.2	86.7	0.7 ⁵
Phe	86.7	86.5	86.0	86.0	86.4	87.1	85.2	86.2	87.2	84.0 [*]	86.8	84.9	86.5	87.2	87.4	87.5	0.6 ⁵
Lys	89.8	89.9	90.4	89.6	89.7	92.9 [*]	89.7	91.2	91.6	90.3	90.9	89.9	90.6	91.6	92.3 [*]	91.4	0.6 ^{5,6}
His	88.7	87.4	87.6	87.4	87.5	89.2	87.7	89.1	89.7	87.6	89.6	86.9	89.2	88.9	90.2	89.1	0.7 ⁵
Arg	91.6	93.0	91.2	92.3	91.2	92.5	90.7	91.4	93.5	89.7	92.7	92.4	91.2	92.2	92.6	93.1	0.7 ^{5,6}
Trp	88.2	89.0	89.3	88.4	88.3	87.4	86.7	88.4	88.1	87.0	87.4	88.5	87.4	89.1	88.0	86.0 [*]	0.5 ^{5,6}
Starch	97.6	99.4 [*]	98.1	98.9 [†]	98.1	98.8	97.0	97.4	97.5	97.3	98.3	96.7	97.2	96.9	97.3	98.3	0.3 ⁵

558 ¹ Reduced nutrient density negative control.

559 ² Nutrient adequate positive control.

560 ³ Non-essential amino acids.

561 ⁴ Essential amino acids.

562 ⁵ Significant effect of diet (P < 0.05). Means in the same column are significantly different from the negative control, *P < 0.05, †P < 0.10.

563 ⁶ Significant effect of block (P < 0.05).

564 **Table 10.** Apparent digestible amino acid intake (g/day) of broiler chicks fed reduced crude protein and amino acid diets and novel proteases
 565 from hatch to 17 days post-hatch (Experiment 2)

Nutrient	NC ¹	Neutral protease								Acid protease						PC ²	SE
		1	2	3	4	5	6	7	8	1	2	3	4	5	6		
NEAA ³																	
Asp	0.68	0.40*	0.68	0.67	0.62	0.68	0.55*	0.63	0.69	0.58*	0.68	0.55*	0.64	0.64	0.67	0.72	0.02 ^{5,6}
Ser	0.33	0.21*	0.34	0.34	0.30	0.35	0.28*	0.32	0.35	0.29*	0.34	0.28*	0.31	0.31	0.33	0.35	0.01 ^{5,6}
Glu	1.31	0.76*	1.28	1.28	1.16*	1.28	1.07*	1.19	1.31	1.13*	1.30	1.10*	1.21	1.21	1.25	1.35	0.04 ^{5,6}
Pro	0.41	0.26*	0.41	0.41	0.36 [†]	0.40	0.33*	0.35*	0.41	0.38	0.39	0.34*	0.37	0.40	0.37	0.41	0.01 ^{5,6}
Gly	0.28	0.16*	0.27	0.27	0.24*	0.28	0.22*	0.26	0.29	0.24*	0.28	0.23*	0.26	0.26	0.27	0.29	0.01 ^{5,6}
Ala	0.34	0.20*	0.33	0.33	0.30*	0.34	0.28*	0.31	0.35	0.30 [†]	0.33	0.29*	0.31	0.31	0.33	0.34	0.01 ^{5,6}
Cys	0.10	0.06*	0.10	0.10	0.08*	0.10	0.08*	0.09	0.10	0.08*	0.09	0.07*	0.09	0.10	0.09	0.09	0.00 ^{5,6}
Tyr	0.23	0.13*	0.23	0.23	0.21*	0.21	0.18*	0.21 [†]	0.23	0.19*	0.23	0.19*	0.21	0.20*	0.22	0.24	0.01 ⁵
EAA ⁴																	
Thr	0.27	0.16*	0.27	0.27	0.25	0.28	0.22*	0.26	0.29	0.24 [†]	0.28	0.23*	0.26	0.26	0.27	0.29	0.01 ^{5,6}
Val	0.31	0.19*	0.31	0.30	0.28	0.31	0.25*	0.29	0.32	0.27*	0.31	0.26*	0.29	0.29	0.30	0.32	0.01 ^{5,6}
Met	0.20	0.13*	0.21	0.21	0.18	0.21	0.17*	0.18 [†]	0.22	0.17*	0.21	0.17*	0.18*	0.23*	0.21	0.23*	0.01 ^{5,6}
Iso	0.29	0.17*	0.28	0.28	0.26 [†]	0.29	0.23*	0.27	0.29	0.25*	0.29	0.23*	0.27	0.27	0.28	0.30	0.01 ^{5,6}
Leu	0.61	0.36*	0.58	0.57	0.53*	0.58	0.49*	0.54 [†]	0.60	0.52*	0.59	0.50*	0.55	0.55	0.57	0.61	0.02 ^{5,6}
Phe	0.36	0.20*	0.33	0.33	0.32 [†]	0.34	0.28*	0.31*	0.34	0.29*	0.34	0.28*	0.32*	0.32 [†]	0.33	0.37	0.01 ^{5,6}
Lys	0.47	0.30*	0.48	0.46	0.45	0.53 [†]	0.40	0.45*	0.53	0.48	0.49	0.43	0.45	0.48	0.52	0.50	0.01 ^{5,6}
His	0.19	0.11*	0.18	0.18	0.16 [†]	0.17	0.15*	0.17	0.19	0.16 [†]	0.18	0.15*	0.17	0.17	0.18	0.19	0.01 ^{5,6}
Arg	0.48	0.28*	0.46	0.47	0.43	0.47	0.38*	0.44	0.50	0.41*	0.47	0.39*	0.44	0.44	0.46	0.51	0.01 ^{5,6}
Trp	0.09	0.06*	0.09	0.09	0.08*	0.08 [†]	0.07*	0.09	0.09	0.08	0.08	0.09	0.08*	0.08	0.08	0.08*	0.00 ^{5,6}

566 ¹ Reduced nutrient density negative control.

567 ² Nutrient adequate positive control.

568 ³ Non-essential amino acids.

569 ⁴ Essential amino acids.

570 ⁵ Significant effect of diet (P < 0.05). Means in the same column are significantly different from the negative control, *P < 0.05, [†]P < 0.10.

571 ⁶ Significant effect of block (P < 0.05).

572 **Table 11.** Growth performance of broilers fed reduced nutrient density diets and novel
 573 proteases from hatch to 18 days post-hatch (Experiment 3)

Diet	Protease	Phytase, FTU/kg	Feed intake, g	BW gain, g	FCR, g:g	Mortality, %
Negative control		500	807.6	508.7	1.573	2.0
	Neutral					
	1	500	724.8	374.5*	1.801*	4.0
	2	500	781.5	486.0	1.620	2.0
	3	500	771.7	483.8	1.616	0.0
	4	500	789.2	797.9	1.613	6.0
	5	500	806.0	512.8	1.575	6.0
	6	500	832.9	517.9	1.617	0.0
	7	500	812.0	498.1	1.645	4.0
	8	500	834.6	522.9	1.609	4.0
	Acid					
	1	500	760.2	476.0	1.609	0.0
	3	500	752.5	460.9	1.646	0.0
	4	500	813.9	471.0	1.739 [†]	0.0
	5	500	789.6	488.0	1.640	2.0
		1500	830.7	529.9	1.577	4.0
		3000	837.0	521.8	1.617	2.0
Positive control		500	879.7	556.3	1.588	2.0
	SE		23.3	17.3	0.04	0.02
	Diet P-value		0.0021	< 0.0001	0.0553	0.59
	Block P-value		0.32	< 0.0001	< 0.0001	0.98

574 Means in the same column are significantly different from the negative control, *P < 0.05, [†]P
 575 < 0.10.

576 **Table 12.** Apparent ileal amino acid and starch digestibility of broiler chicks fed reduced crude protein and amino acid diets and novel proteases
 577 from hatch to 18 days post-hatch (Experiment 3)

Nutrient	NC ¹	Neutral protease								Acid protease				Phytase, FTU		PC ²	SEM
		1	2	3	4	5	6	7	8	1	3	4	5	1500	3000		
NEAA ³																	
Asp	84.2	83.9	83.2	82.8	82.7	83.2	83.4	82.2	81.1*	83.5	82.7	84.2	83.2	83.0	83.2	85.2	0.6 ⁵
Ser	85.0	87.4 [†]	84.2	82.4*	84.5	83.8	83.7	82.0*	80.6*	84.4	84.0	84.0	83.7	83.1	83.3	84.6	0.6 ⁵
Glu	92.2	92.3	92.0	91.8	92.1	92.1	91.9	91.1	90.6*	92.5	92.0	92.3	92.3	91.5	91.8	91.8	0.3 ⁵
Pro	89.8	91.7	89.0	88.5	88.9	89.1	89.6	88.3	88.5	89.6	89.1	89.1	90.0	89.2	89.4	90.1	0.6
Gly	83.0	85.3	81.1	81.4	80.6	81.4	82.0	80.4 [†]	78.8*	81.5	81.7	81.8	82.5	81.0	81.6	83.7	0.6 ⁵
Ala	83.9	82.6	82.3	82.0	81.5	82.4	83.3	81.6	80.7*	82.9	82.6	82.8	83.7	82.1	82.5	84.7	0.7 ⁵
Cys	80.1	77.0	76.5	77.0	76.4	77.3	77.8	76.2	75.2	77.9	77.7	78.1	79.0	78.7	77.4	77.2	1.1 ⁶
Tyr	83.8	88.2	85.0	85.5	86.2	85.8	86.8	85.0	80.8	84.5	83.4	85.7	86.0	82.9	83.8	86.2	1.4 ^{5,6}
EAA ⁴																	
Thr	84.0	86.2	82.6	81.3*	82.5	83.0	83.9	81.6 [†]	81.3*	82.8	83.0	83.4	83.5	82.5	83.1	84.3	0.6 ⁵
Val	84.3	86.6	83.0	82.8	82.3	82.9	83.6	82.4	81.7 [†]	83.1	83.1	84.1	84.4	82.8	82.7	85.2	0.7 ⁵
Met	94.1	95.1	93.8	93.3	94.1	94.1	94.7	94.0	93.5	94.2	94.1	94.5	94.4	93.7	94.5	94.8	0.4 ⁶
Iso	85.4	88.0*	84.6	84.3	84.2	84.8	84.8	84.6	82.6*	85.4	84.5	85.5	85.5	84.5	84.5	86.1	0.6 ^{5,6}
Leu	86.8	89.9*	86.2	86.2	85.9	86.4	86.4	85.4	84.1*	86.9	86.0	86.8	86.9	85.7	86.3	87.1	0.6 ^{5,6}
Phe	87.7	89.7	85.7	85.9	86.0	86.7	87.3	85.7	84.4*	86.8	84.9	87.0	86.8	86.0	86.5	86.9	0.8 ^{5,6}
Lys	91.1	92.7*	90.5	90.0	89.8	91.1	90.7	90.1	89.6	90.9	90.6	90.6	91.2	90.1	90.9	91.3	0.4 ⁵
His	88.0	90.4*	86.5	86.4	86.8	87.6	88.1	86.5	85.8 [†]	86.7	87.2	87.8	88.7	86.6	87.0	88.5	0.6 ⁵
Arg	89.0	88.8	87.8	87.3	86.0*	88.2	88.9	87.5	86.4*	88.4	88.1	88.9	89.1	87.8	88.5	89.8	0.5 ⁵
Starch	82.4	71.9	78.4	83.6	80.1	80.0	83.0	76.0	79.2	82.5	79.4	86.6	86.6	85.1	82.6	76.1	4.6 ⁶

578 ¹ Reduced nutrient density negative control.

579 ² Nutrient adequate positive control.

580 ³ Non-essential amino acids.

581 ⁴ Essential amino acids.

582 ⁵ Significant effect of diet (P < 0.05). Means in the same column are significantly different from the negative control, *P < 0.05, [†]P < 0.10.

583 ⁶ Significant effect of block (P < 0.05).

584 **Table 13.** Apparent digestible amino acid intake (g/day) of broiler chicks fed reduced crude protein and amino acid diets and novel proteases
 585 from hatch to 18 days post-hatch (Experiment 3)

Nutrient	NC ¹	Neutral protease								Acid protease				Phytase, FTU		PC ²	SEM
		1	2	3	4	5	6	7	8	1	3	4	5	1500	3000		
NEAA ³																	
Asp	0.64	0.55*	0.57	0.54*	0.57	0.59	0.64	0.60	0.59	0.55*	0.52*	0.64	0.56*	0.62	0.60	1.04*	0.02 ⁵
Ser	0.35	0.33	0.33	0.29*	0.34	0.32	0.35	0.32	0.31	0.32	0.29*	0.34	0.30*	0.33	0.33	0.50*	0.01 ⁵
Glu	1.62	1.43 [†]	1.51	1.47	1.56	1.58	1.65	1.56	1.55	1.49	1.43*	1.63	1.51	1.59	1.61	2.16*	0.05 ⁵
Pro	0.47	0.40*	0.42*	0.41*	0.44	0.47	0.51	0.45	0.49	0.41*	0.43	0.48	0.43	0.50	0.48	0.72*	0.01 ^{5,6}
Gly	0.29	0.26	0.25 [†]	0.25*	0.25 [†]	0.26	0.28	0.27	0.26	0.24*	0.24*	0.28	0.26	0.28	0.27	0.40*	0.01 ⁵
Ala	0.28	0.24*	0.25	0.24*	0.24*	0.25	0.29	0.27	0.27	0.24*	0.24*	0.28	0.25	0.27	0.26	0.41*	0.01 ^{5,6}
Cys	0.12	0.10*	0.10*	0.10*	0.10*	0.11	0.11	0.11	0.11	0.10*	0.10*	0.11	0.10*	0.12	0.11	0.15*	0.00 ^{5,6}
Tyr	0.18	0.18	0.18	0.18	0.19	0.17	0.21*	0.17	0.17	0.17	0.15*	0.18	0.18	0.17	0.18	0.26*	0.01 ⁵
EAA ⁴																	
Thr	0.30	0.27*	0.28	0.25*	0.28	0.29	0.33	0.29	0.30	0.27 [†]	0.26*	0.31	0.29	0.31	0.30	0.42*	0.01 ^{5,6}
Val	0.30	0.27*	0.27	0.26*	0.27	0.27	0.31	0.29	0.29	0.26*	0.25*	0.31	0.29	0.30	0.28	0.46*	0.01 ⁵
Met	0.22	0.19	0.22	0.19	0.23	0.23	0.27*	0.24 [†]	0.24 [†]	0.21	0.21	0.26*	0.24	0.23	0.26*	0.36*	0.01 ⁵
Iso	0.28	0.25	0.25	0.23*	0.25	0.25	0.29	0.27	0.26	0.25	0.23*	0.28	0.26	0.27	0.26	0.43*	0.01 ⁵
Leu	0.50	0.46	0.47	0.45 [†]	0.47	0.47	0.52	0.49	0.47	0.45	0.43*	0.51	0.46	0.49	0.49	0.74*	0.01 ⁵
Phe	0.35	0.31*	0.31*	0.29*	0.31	0.33	0.37	0.33	0.33	0.30*	0.29*	0.35	0.32	0.34	0.34	0.50*	0.01 ⁵
Lys	0.52	0.45*	0.48	0.44*	0.46 [†]	0.52	0.53	0.51	0.52	0.46 [†]	0.44*	0.53	0.48	0.50	0.52	0.75*	0.02 ⁵
His	0.18	0.16*	0.16*	0.15*	0.16	0.17	0.19	0.18	0.18	0.15*	0.16*	0.18	0.18	0.18	0.18	0.27*	0.01 ⁵
Arg	0.45	0.39*	0.40 [†]	0.37*	0.38*	0.42	0.48	0.44	0.43	0.38*	0.38*	0.46	0.41	0.45	0.44	0.71*	0.01 ⁵

586 ¹ Reduced nutrient density negative control.

587 ² Nutrient adequate positive control.

588 ³ Non-essential amino acids.

589 ⁴ Essential amino acids.

590 ⁵ Significant effect of diet (P < 0.05). Means in the same column are significantly different from the negative control, *P < 0.05, [†]P < 0.10.

591 ⁶ Significant effect of block (P < 0.05).