

Comparison of potato (*Solanum tuberosum* L.) tuber size distribution fitting methods and evaluation of the relationship between soil properties and estimated distribution parameters

by Mhango, J.K., Hartley, W., Harris, W.E. and Monaghan, J.M.

Copyright, publisher and additional information: Publishers' version distributed under the terms of the [Creative Commons Attribution License](#)

[DOI link to the version of record on the publisher's site](#)



Mhango, J.K., Hartley, W., Harris, W.E. and Monaghan, J.M. (2021) 'Comparison of potato (*Solanum tuberosum* L.) tuber size distribution fitting methods and evaluation of the relationship between soil properties and estimated distribution parameters', *The Journal of Agricultural Science*.

15 December 2021

Crops and Soils Research Paper

Cite this article: Mhango JK, Hartley W, Harris WE, Monaghan JM (2021). Comparison of potato (*Solanum tuberosum* L.) tuber size distribution fitting methods and evaluation of the relationship between soil properties and estimated distribution parameters. *The Journal of Agricultural Science* 1–15. <https://doi.org/10.1017/S0021859621000952>

Received: 5 May 2021

Revised: 1 November 2021

Accepted: 12 November 2021

Keywords:


Nitrogen; phosphorous; potato yield prediction; precision farming; tuber number prediction

Author for correspondence:

J. M. Monaghan,

E-mail: jmmonaghan@harper-adams.ac.uk

Comparison of potato (*Solanum tuberosum* L.) tuber size distribution fitting methods and evaluation of the relationship between soil properties and estimated distribution parameters

J. K. Mhango, W. Hartley, W. E. Harris and J. M. Monaghan 

Agriculture and Environment Department, Harper Adams University, TF10 8NB, Edgmond, Shropshire, UK

Abstract

Accurate estimation of tuber size distribution (TSD) parameters in discretely categorized potato (*Solanum tuberosum* L.) yield samples is desirable for estimating modal tuber sizes, which is fundamental to yield prediction. In the current work, systematic yield digs were conducted on five commercial fields ($N = 119$) to compare the Weibull, Gamma and Gaussian distribution functions for relative-likelihood-based goodness-of-fit to the observed discrete distributions. Parameters were estimated using maximum likelihood estimation (MLE) for the three distributions but were also derived using the percentiles approach for the Weibull distribution to compare accuracy against the MLE approaches. The relationship between TSD and soil nutrient variability was examined using the best-fitting model's parameters. The percentiles approach had lower overall relative likelihood than the MLE approaches across five locations, but had consistently lower Root Mean Square Error in the marketable tuber size range. Negative relationships were observed between the percentile-based shape parameter and the concentrations of Phosphorus and Nitrogen, with significant (non-zero-overlapping 95% confidence interval) regression coefficients for P (-0.74 ± 0.33 for distribution of proportional tuber numbers and -1.3 ± 0.62 for tuber weights). Stem density was negatively associated with the scale and mode of tuber number (regression coefficients -0.98 ± 0.63 and -1.08 ± 0.78 respectively) and tuber weight (regression coefficients -0.99 ± 0.78 and -1.04 ± 0.69 respectively) distributions. Phosphorus is negatively related to the scale of the tuber-number-based distribution while positively associating with the tuber weight distribution. The results suggest that excess P application was associated with the increase in small tubers that did not contribute significant weight to the final yield.

Introduction

Potato (*Solanum tuberosum* L.) production is a high input, high output operation driven by well-developed, technologically advanced markets which depend on a consistent supply of high-quality potatoes. Nevertheless, high output alone does not necessarily translate into high returns for growers due to the selectivity of processing factories for tuber size grades (Machakaire *et al.*, 2016). This occurs especially in the pre-fried potato processing sector which accounts for 62% of the global processed potato market (Keijbets, 2008). Agronomic techniques for optimizing tuber size distribution (TSD) are therefore an important consideration for potato growers and researchers.

Several methods of quantitatively describing TSD have been proposed in the literature. Travis (1987) described TSD using the spread of tuber sizes around the modal grade assuming a Gaussian distribution, allowing the determination of a coefficient of variation (CV) as an index for TSD. Ideally, farmers can use this to conduct mid-season TSD assessments, which can then support management decisions on vine desiccation timing. Struik *et al.* (1990) and others subsequently supported the Travis (1987) method. However, the probability densities plotted by Struik *et al.* (1991) revealed that the TSD by weight skews to the right and the Gaussian distribution may not necessarily capture the spread of the data.

Several alternatives have been suggested like the Weibull (Nemecek *et al.*, 1996; Bussan *et al.*, 2007) Log-normal (Marshall *et al.*, 1993) and the Gamma (Aliche *et al.*, 2019) functions. Whilst these alternative functions fit TSD more accurately, their parameters are often determined using maximum likelihood estimation (MLE), which makes it non-ideal for quick field assessment by non-statisticians. A Gaussian distribution is therefore often assumed due to the simplicity with which its parameters (i.e. mean and standard deviation) can be determined from yield digs. Currently, TSD is mostly evaluated using the percentage of marketable tuber weight in the total tuber yield.

© The Author(s), 2021. Published by Cambridge University Press. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

CAMBRIDGE
UNIVERSITY PRESS

Potato growers, and subsequently researchers, target different tuber sizes at harvest depending on localized outlet market demands (Wurr *et al.*, 1993; Taylor *et al.*, 2018), making it difficult to standardize the size classes and objectively compare TSD from different studies. A positive association between soil Nitrogen (N) concentration and TSD has been consistently reported by several authors over the past five decades, mostly when TSD is measured as the proportion or absolute quantity of yield above a weight or transversal diameter threshold (Schippers, 1968; Porter and Sisson, 1991; Arsenault *et al.*, 2001; Gao *et al.*, 2018). The overall effect of N on the shape, location and scale of the TSD has not been consistently demonstrated partly due to the lack of standardized and generally accepted parameters for measurement (Wurr *et al.*, 1993).

The influence of Potassium (K) fertilization on potato yield and TSD is also widely studied (Dickins *et al.*, 1962; Birch *et al.*, 1967; Allison *et al.*, 2001) and soil K replacement based on crop removal is generally accepted as a management strategy in commercial production systems. For Phosphorus (P), Rosen and Bierman (2008) found that incremental rates between 0 and 74 kg/ha did not affect the percentage of marketable (>85 g) Russet Burbank yield in loamy sand soil with medium to high P concentrations (25–33 mg/kg Bray P1). This was attributed to an increase in the number of small tubers happening concurrently with a reduction in the percentage of large tubers as P rate increased. Quoting Westermann and Kleinkopf (1985), Rosen and Bierman (2008) attribute this response to a shift in dry matter partitioning from tubers to vegetative growth as leaf P increased. The response of TSD to P can be affected by P interactions with other elements in the soil.

Phosphorus is known to exhibit antagonistic relationships with zinc (Zn) and magnesium (Mg) under alkaline conditions and iron (Fe) and Aluminium (Al) under strongly acidic conditions (Rietra *et al.*, 2017). These four elements precipitate P out of the soil solution and render it unavailable for plant uptake,

hence confounding the effect of P fertilization on agronomic parameters. The influence of Mg is worth consideration because it is often applied in potato fields as Epsom salt to control Mg-deficiency-related leaf chlorosis. Finally, the role of the Sulphate counter ion in $MgSO_4$ salts and K_2SO_4 , as well as the inherent soil variability in Sulphur (S) is rarely studied and is often discounted subject of speculation (Simmons and Kelling, 1987). However, Caldiz *et al.* (2018) found that 61% of the variation in the proportion of small tubers (<50 mm) was explained by the variation in soil S content ($P < 0.05$). An increase in S concentration in the soil had a positive correlation with the percentage of small tubers in the final yield. Within-field variation in S may ultimately be correlated to variation in TSD and evidence from Caldiz *et al.* (2018) supports the hypothesis of an increase in small-sized tubers observed in K_2SO_4 applications by Henderson (1965).

The objectives of the current study were to (a) identify which of the Weibull, Gamma and Gaussian distribution functions was the best at simulating observed TSD parameters, and; (b) to utilize the best-fitted distribution function to study the influence of soil macronutrient availability on TSD.

Methodology

Site characterization

The study was conducted at four sites as summarized in Table 1. Deaton 6 and HF7 sites were located in the East of England (Lincolnshire) on reclaimed marshland with a shallow water table and high organic matter content. There was variation in soil physical and chemical properties across the field due to the presence of Roddons, historical features in drained marshland soils where silty clay soils follow the course of historical streams and waterways. Horse Foxhole and Buttery Hill were located in the West of England (Shropshire) on well-drained slightly stony,

Table 1. Summarized information of the study sites

Field Name	Year	Coordinates	Variety	N ¹	Fertilizer ²	Nutrient Rate (kg/ha)
HF7	2020	53°12'40.71"N 0°24'49.76"W	Maris Piper	30	Nitrogen	107
					Phosphorus	94
					Potassium	290
B.Hill ³	2020	52°46'22.05"N 2°25'40.46"W	Amora	30	Nitrogen	173
					Potassium	94
Deaton 6	2019	53°12'20.97"N 0°21'55.06"W	Maris Piper	12	Nitrogen	100
					Phosphorus	150
					Potassium	307
C.Leasow ⁴	2020	52°46'15.21"N 2°21'37.57"W	P.Dell ⁵	18	Nitrogen	167
					Potassium	89
H.Fxhole ⁶	2019	52°46'26.94"N 2°25'49.38"W	Amora	29	Nitrogen	125
					Phosphorus	140
					Potassium	238
					Magnesium	40
					Sulphur	75

1 = Number of samples collected. 2 = Main fertilizer forms were N for Nitrogen, P₂O₅ for Phosphorus, K₂O for Potassium, SO₃ for Sulphur and MgSO₄·7H₂O for Sulphur and Magnesium. 3 = Buttery Hill. 4 = Crabtree Leasow. 5 = Pentland Dell. 6 = Horse Foxhole.

sandy loam soil subtended by weathered sandstone with low variation in soil nutrients across the field. Additional tuber sampling for TSD modelling (without soil analysis) was conducted at Crabtree Leasow, also located in the West of England with similar soil and weather conditions to Buttery Hill and Horse Foxhole. Thus, our study sites are representative of a broad range of soil conditions as is typical in potato production agriculture. Planting and field management was carried out by the respective farmer at each field. Consequently, land preparation was conducted similarly in all fields by ploughing at 30 cm depth followed by bed-forming at 90 cm between rows and destoning. Fertilizer was applied uniformly by broadcasting macro and micronutrients based on soil analysis as summarized in Table 1. Deaton 6 and HF7 were irrigated by drip irrigation while a hose reel irrigator was used at Horse Foxhole and Buttery Hill. All management practices were conducted uniformly across the field throughout the growing period.

Sampling design

A field survey was designed at each of the five fields using a model-based sampling approach to determine representative soil sampling locations that captured the variability in the field. The number of samples collected at all locations are as provided in Table 1. Soil sampling was done at all sites except Crabtree Leasow (thus not included in the soil analysis). Accordingly, the soil samples were collected at Buttery Hill ($N=23$), HF7 ($N=24$), Deaton 6 ($N=12$) and Horse Foxhole ($N=23$). Soil macronutrient quantities are known to correlate with organic matter content (Yang *et al.*, 2011), which in turn influences soil colour (Costa *et al.*, 2020). The Soil Brightness Index (SBI) as described by Mponela *et al.* (2020), was chosen as a substitute for spatially modelling the soil colour differences at each field and generating strata for a stratified random sampling design. At each field, the SBI was calculated at least 1 day per month for 3 months prior to crop emergence and then the average SBI was calculated. The SBI at each site was calculated using atmospherically corrected (Level-2A) satellite imagery of 10 m resolution acquired by the Sentinel-2 satellite, on manually inspected cloud-free days. Each field was delineated into three clusters by k -means clustering ($k=3$) based on SBI to generate zones of relative homogeneity which formed the sampling strata. A sampling unit of 6 m by 6 m (36 m^2) was chosen to cover the accuracy specification of the GarminTM eTrex 20 GPS receiver that was used for soil sampling. A grid of 36 m^2 quadrats was imposed across a rasterized SBI surface then random quadrats were drawn from each stratum.

Power analysis to determine the sample size for the survey was calculated to resolve SBI variability with a statistical power of 0.8 as recommended by Cohen (1988). The effect size was estimated based on the expected within-field contrast between the k -means cluster with the lowest SBI value (dark soil) and the one with the highest SBI value (bright soil) in the sample. The effect size was therefore calculated as the difference between the mean SBI of dark soils and the combined mean of the medium and light clusters divided by the standard deviation of the entire dataset. R v4.0.2 (R Core Team, 2019) was used to calculate the sample size using the 'pwr.t.test' function from the 'pwr' package (Champely *et al.*, 2018). Once the sample size of each cluster was determined, sampling locations were selected by randomly selecting quadrats from the grid imposed on the SBI raster. The quadrats were georeferenced and assigned with unique identifier codes then exported as a GPX file into the GPS receiver for

tracking during the soil and yield sampling. All raster analysis steps were performed using ArcGIS (ESRI, 2018).

Soil and nutrient analysis

Soil sampling was conducted before planting, but after fertilizer incorporation. Soils were sampled using a 30 cm auger. Samples were collected in triplicate from each quadrat and mixed to form a composite sample. The soil samples were then air-dried at 30°C for 72 h. After air drying, the soils were ground and sieved (<2 mm) prior to analysis. The percentages of sand, silt and clay in each soil sample were determined using the sedimentation method (Jackson *et al.*, 1986). Hydrogen peroxide was used to oxidize organic matter, after which the particle size distribution was determined through sieving and sedimentation.

Soil samples were analysed for N, C and S using the Dumas method (Kirsten and Hesselius, 1983). Air-dried soils (0.25 g for N and 0.15 g for C and S) were passed through a furnace at 1000°C in the presence of oxygen. The oxidized gases were then detected and measured using a thermal conductivity cell. The Olsen method (Page, 1982) was used to estimate available P in the soil. Sodium bicarbonate was used to extract P from the soil into solution and form phosphomolybdate after reaction with ammonium molybdate. The phosphomolybdate was reduced by ascorbic acid to form a blue complex whose concentration was measured spectrophotometrically at 880 nm. Concentrations of K and Mg were determined by flame photometry using ammonium nitrate as an extractant as described in (Jackson *et al.*, 1986).

Yield data collection

At every sampling location, the number of plants within a one-metre row was counted and recorded as a measure of plant density. At harvest, all the plants in the one-metre length were carefully uprooted with a spade and the number of main stems was counted as a measure of stem density. The excavation was carried out carefully to minimize any loss of tubers. All tubers were separated from their stolon and stored for further processing.

The number of tubers at each sampling point was counted. At all locations except for Horse Foxhole, all tubers with a transversal diameter greater than or equal to 25 mm were graded using potato sizing squares into 10 mm size grades up to 65 mm, with all tubers over 65 mm in diameter placed in one bin. All tubers under 25 mm diameter were binned into one grade. After grading, the tuber number in each category was counted and its weight was determined at 0.01 g accuracy. For Horse Foxhole, the tubers were separated into typical commercial grading of 0–25 mm, 25 mm–45 mm, 45–65 mm and greater than 65 mm.

Data analysis

TSD was modelled using the Gaussian (Travis, 1987; Aliche *et al.*, 2019), Gamma (Aliche *et al.*, 2019) and Weibull (Nemecek *et al.*, 1996; Bussan *et al.*, 2007) distributions. The Gaussian distribution is the most widely adopted of the three distributions, with the mode equalling the mean of the distribution and its parameter estimation is simple and intuitive as described in Travis (1987). For the Gamma distribution, the probability density is as described in Aliche *et al.* (2019), with respect to TSD in potatoes. The Weibull distribution density function was described by Nemecek *et al.* (1996). The Gamma and Weibull distributions are considered to be flexible to right-skewed data, common in

potato TSD (Nemecek *et al.*, 1996; Bussan *et al.*, 2007; Aliche *et al.*, 2019), and hence they can potentially more accurately estimate modal tuber size than the Gaussian distribution. Love and Thompson-Johns (1999) demonstrated that agronomic practices such as wider plant spacing can also lead to left-skewed TSD, making the Weibull distribution an ideal candidate for modelling the shape of the distribution flexibly. Figure 1 illustrates the effect of changing the shape parameter on a conceptual Weibull probability density curve. Generally, the distribution resolves to an approximately symmetrical distribution with a shape value of 3.6 and approaches right-skewness or left-skewness above or below that respectively (Lai *et al.*, 2006). Thus, the Weibull distribution is ideal in this application to model both left and right-skewed data.

The MLE approach was used to obtain estimates for the mean and standard deviation of the Gaussian distribution, and the scale and shape parameters for the Weibull and Gamma distributions. The 'fitdstrplus' R package (Delignette-Muller and Dutang, 2015) was used to obtain the parameter estimations for all three distributions from the right-censored interval potato TSD data of the current study. Accordingly, the likelihood of each parameter θ in each distribution was fitted in Eqn (1):

$$L(\theta) = \prod_{j=1}^{N_{\text{left C}}} F(x_j^{\text{upper}}|\theta) \times \prod_{k=1}^{N_{\text{right C}}} 1 - F(x_k^{\text{lower}}|\theta) \times \prod_{m=1}^{N_{\text{int C}}} (F(x_m^{\text{upper}}|\theta) - F(x_j^{\text{lower}}|\theta)) \quad (1)$$

Using the cumulative distributions (F), where x_j^{upper} represented the upper values defining the $N^{\text{left C}}$ left-censored observations, x_k^{lower} represented the lower values defining the $N^{\text{right C}}$ right-censored observations, x_m^{lower} and x_m^{upper} represented the intervals defining the $N^{\text{int C}}$ interval-censored observations (Delignette-Muller and Dutang, 2015).

The Weibull distribution parameters were also estimated using the percentiles method (Dubey, 1967). The motivation for including this method was the simple closed-form procedure for calculating parameter estimates, which potentially makes it easier for non-statisticians to calculate the parameters and make an inference. The adoption of the Weibull MLE approach in TSD research is very low, with recent studies still using Gaussian mean estimates (Buhrig *et al.*, 2015) or simply relative frequencies (Andrade *et al.*, 2021; van Dijk *et al.*, 2021) to describe TSD and its changes with respect to agronomic treatments. This is attributable to non-familiarity with the approach due to the ubiquity of least-squares-based procedures and simplicity of Gaussian parameter estimates, which still provide useful models as reported in Travis (1987). Estimating TSD using the Weibull percentiles approach provides a simpler alternative to the poorly adopted MLE approach, contingent upon achieving a comparable empirical accuracy between the two approaches, hence their comparison in this study. Accordingly, the shape parameter ($\hat{\beta}_{\text{Weibull}}$) of the Weibull distribution for the percentiles method was estimated by linearizing the cumulative distribution function at two different discrete diameters (e.g. 45 and 65 mm) then combining the two equations to solve for $\hat{\beta}_{\text{Weibull}}$ as in Eqn (2):

$$\hat{\beta}_{\text{Weibull}} = (\ln x_i - \ln x_j)^{-1} \times \left\{ \ln \left[\frac{\ln(1 - f(x_j))}{\ln(1 - f(x_i))} \right] \right\} \quad (2)$$

The scale parameter $\hat{\alpha}_{\text{Weibull}}$ was then calculated using $\hat{\beta}_{\text{Weibull}}$ and the cumulative density at one known quantile as in Eqn (3):

$$\hat{\alpha}_{\text{Weibull}} = \frac{x_i}{[-\ln(1 - f(x_i))]^{\frac{1}{\hat{\beta}_{\text{Weibull}}}}} \quad (3)$$

where $f(x_{i..j})$ represented the cumulative probability of tuber number or weight at x_i or x_j tuber diameter and $x_{i..j}$ where the chosen i^{th} or j^{th} discrete diameter of a tuber

In each sample, the estimated parameters of the four distribution-fitting approaches (Gaussian, Gamma, Weibull with MLE and Weibull with percentiles approach) were used to estimate the tuber numbers in each original discrete size grade, creating fully specified ordered discrete distributions. The estimation was done by multiplying the estimated probability of each size grade (the difference between the upper limit and lower limit cumulative probabilities of each size grade) by the observed total tuber number. The Weibull percentiles approach was fitted at the percentiles corresponding to 45 and 65 mm tuber sizes, selected because this represents the marketable range for main-crop potatoes. In this case, the percentiles were the cumulative proportions of the tuber number or weight at 45 or 65 mm, and were therefore different for every sample. The logarithm of the relative likelihood (hereinafter referred to as log relative likelihood) of the predicted discrete distribution (relative to the actual discrete distribution) was then calculated as illustrated in Lindsey (1974). The likelihood of the discrete distribution was calculated as the likelihood of a fully specified multinomial distribution in Eqn (4), where P is the probability of the j^{th} category and n is the frequency of the j^{th} category.

$$L(P) = \prod_j P_j^{n_j} \quad (4)$$

The log relative likelihood was then used as a ranking index of the plausibility of the models, with the most plausible model having the highest likelihood. The log relative likelihood was calculated as in Eqn (5):

$$L_{RR} = \sum_j^n [\ln(\hat{P}_j) - \ln(P_j)] \quad (5)$$

where n_j is the observed frequency of a category, \hat{P}_j is the probability of the predicted distribution and P_j is the likelihood of the observed distribution.

Uncertainty in the log relative likelihood was assessed using 95% confidence intervals. Fisher's information was used to estimate the variability of the estimate. The Fisher's information was derived by negating the expected value of the second derivative of the log-likelihood equation (Ly *et al.*, 2017), which for the multinomial distribution used was defined as in Eqn (6):

$$I(\theta_j) = -E\left(-\frac{n_j}{P_j^2}\right) \quad (6)$$

where θ_j is the log relative likelihood of the j^{th} category, n_j is the frequency of the j^{th} category and P_j is the probability of the j^{th} category. The reciprocal of the square of the fisher's information was used as the standard error component of the confidence interval formula.




Fig. 1. The effect of changing the shape parameter on a conceptual Weibull probability density curve.

The main purpose of fitting the theoretical continuous distributions to the discretely measured tuber size fractions is to maximize the accuracy of the estimate of the modal tuber size grade of the distribution, which is the marketable component of the production. This falls within the 45–65 mm size bands for typical maincrop varieties. Therefore, the main purpose of the modelling presented here was to determine which of the distributions-fitting approaches maximized the likelihood of this tuber size fraction, using the observed tuber numbers and weights for this size grade. The log relative likelihood estimates of the 45–65 mm size band were therefore compared for the four models. Additionally, the Root Mean Square Error (RMSE) of prediction for the frequencies or weights in the 45–65 mm size fraction were compared. The RMSE was calculated as in Eqn (7):

$$RMSE = \sqrt{\frac{1}{N} \sum_{i=1}^N (y_i - \hat{y}_i)^2} \quad (7)$$

where N is the number of observations, y_i is the observed value and \hat{y}_i is the predicted value. The modal tuber size of the most plausible model within the 45–65 mm size fraction was considered the best estimate of the distribution’s model. For comparison purposes, the most plausible distribution fitted using the benchmark maximum likelihood approach was used to compare the modal tuber size predictions of the other models using RMSE. The RMSE and log relative likelihood were sample-specific, providing measures of the goodness of fit of the parameter estimates to the observed TSDs without generalizing uncertainties expected at out-of-sample observations. The TSD parameters are only valid for a collected sample, but within-sample hold-out methods (holding out a random size profile within the sample as a test dataset during model fitting) were considered impractical in this case as they would change the TSD of the training set. The mode of the Gaussian distribution was considered to equal the

mean. The mode for the Gamma distribution was calculated in Eqn (8) and the mode for the Weibull distribution was calculated in Eqn (9), as follows:

$$\widehat{Mo}_{Gamma} = \hat{\gamma}_{Gamma} \times (\hat{\alpha}_{Gamma} - 1) \quad (8)$$

$$\widehat{Mo}_{Weibull} = \hat{\alpha}_{Weibull} \times \left(1 - \frac{1}{\hat{\beta}_{Weibull}}\right)^{\frac{1}{\hat{\beta}_{Weibull}}} \quad (9)$$

where \widehat{Mo}_{Gamma} is the estimate of the mode of the Gamma distribution, $\hat{\gamma}_{Gamma}$ is the scale parameter of the Gamma distribution, $\hat{\alpha}_{Gamma}$ is the shape parameter of the Gamma distribution, $\widehat{Mo}_{Weibull}$ is the estimate of the mode of the Weibull distribution, $\hat{\alpha}_{Weibull}$ is the scale of the Weibull distribution and $\hat{\beta}_{Weibull}$ is the shape parameter of the Weibull distribution.

Regression analysis

To assess the significance of the responses of TSD to soil nutrients, linear mixed effect models were computed for the primary macronutrients. The outcome variables of the linear models were the shape and scale parameters of the best-fitting models, treated as new observations. The estimates of the shape and scale parameter at each sample have unique values of uncontrolled uncertainty, contingent upon random error sources and the efficacy of the parameter estimation method. This has the potential of biasing regression coefficients if systematic patterns occur (e.g. spatial autocorrelation in the uncertainty). In this study, the uncertainty was assumed random, with any spatial patterns assumed to be accounted for in subsequent spatial mixed effect covariance structures. Equally, uncertainties in the parameters related to the sampling process leading to varying sample characteristics were considered random, therefore all estimates received the same weight in any further data analysis. The study

locations were also used as the source of random variation. Cognisant of the spatial non-independence of the observations within each location, the mixed effect models included a Matern covariance structure to account for spatial autocorrelation (Stein, 1999; Minasny and McBratney, 2005). The Matern covariance was fitted by restricted error maximum likelihood and all parameters were estimated from the data. Statistical analysis was conducted in R v4.0.2 (R Core Team, 2019) and the spatial regression model was conducted using the SpaMM R package (Rousset and Ferdy, 2014). Statistical significance was evaluated using confidence intervals and the goodness of fit for multivariable regressions was evaluated using nRMSE, which was computed by dividing the RMSE by the mean of the observed variable.

Results

Summary statistics

Summary statistics for the soil and plant variables measured in the study are presented in Table 2. Soil texture ranged from

predominantly sandy silt loams at HF7, through sandy loams at Buttery Hill and Horse Foxhole to predominantly Silty Clays at Deaton 6. Mean total C was highest at Deaton 6 (124.15 g/kg) and HF7 (101.77 g/kg), a reflection of the high organic matter content, while the carbon content of Buttery Hill, and Horse Foxhole were lower. Deaton 6 and HF7 soils also contained higher concentrations of N and S but had lower concentrations of P than the sandy loams. Plant spacing was consistent at approximately 2.5 plants/m² across locations except Horse Foxhole where inconsistent planting spacing led to an average of 5 plants/m². Horse Foxhole also recorded the highest number of stems, tubers and yield per square metre.

The summary statistics of the TSD parameters at the five study sites where the TSD modelling was conducted are presented in Table 3. Mu, the tuber size with the largest probability (modal tuber size with respect to the probability densities of tuber number as determined using the best-fitting TSD model) ranged from 37 mm at Buttery Hill to 51 mm at Horse Foxhole. The same pattern was also reflected in Mu with respect to tuber weight (49 mm at Buttery Hill and 61 mm at Horse Foxhole). Similarly, the scale

Table 2. Summary statistics of key soil and plant variables measured at each study site

Soil.Prop ¹	Deaton 6 Mean (CV ²)	HF7 Mean (CV)	Buttery Hill Mean (CV)	Horse Foxhole Mean (CV)
Clay (%)	38.5 (0.19)	18.3 (0.58)	9.3 (0.15)	14.3 (0.29)
Silt (%)	55.4 (0.12)	38.1 (0.20)	24.6 (0.21)	17.4 (0.30)
Sand (%)	6.1 (0.31)	43.5 (0.30)	66.1 (0.09)	68.3 (0.10)
N (g/kg)	9.4 (0.29)	8.4 (0.45)	1.3 (0.12)	1.7 (0.14)
C (g/kg)	124.2 (0.36)	101.8 (0.46)	12.9 (0.12)	16.3 (0.09)
S (g/kg)	3.3 (0.47)	2.2 (0.51)	0.3 (0.12)	0.3 (0.19)
P (mg/kg)	41.2 (0.15)	42.4 (0.17)	100.0 (0.13)	91.1 (0.12)
pH	7.5 (0.08)	6.6 (0.05)	7 (0.03)	6.6 (0.03)
K (mg/kg)	291.3 (0.27)	272.7 (0.24)	276.8 (0.18)	202.4 (0.19)
Mg (mg/kg)	185.0 (0.29)	80.1 (0.25)	87.4 (0.12)	88.4 (0.14)
Plants/ m ²	2.5 (0.12)	2.5 (0.24)	2.8 (0.26)	5.2 (0.30)
Stems/m ²	9.8 (0.17)	12.4 (0.34)	13.9 (0.28)	17.0 (0.16)
Tubers/m ²	38.7 (0.10)	41.1 (0.28)	35.4 (0.29)	48.3 (0.14)
Yield (kg/m ²)	4.2 (0.12)	5.2 (0.19)	3.4 (0.27)	5.5 (0.11)

1 = Soil Property. 2 = Coefficient of variation.

Table 3. Summary statistics of the TSD parameters at five different sites

Parameter	Deaton 6 Mean (CV ¹)	HF7 Mean (CV)	B.Hill Mean (CV)	H.Foxhole Mean (CV)	C.Leasow Mean (CV)
Mu ² (mm)	49.46 (0.03)	44.50 (0.07)	36.50 (0.08)	51.10 (0.03)	43.70 (0.05)
Mu _{wt} ³ (mm)	56.71 (0.05)	53.57 (0.06)	48.45 (0.07)	61.09 (0.03)	49.49 (0.05)
W.Shape ⁴	5.0 (0.12)	4.5 (0.16)	3.4 (0.23)	4.7 (0.12)	5.15 (0.28)
W.Scale ⁵	51.8 (0.03)	48.8 (0.07)	40.6 (0.07)	55.8 (0.03)	45.93 (0.06)
W.Shape _{wt} ⁶	7.59 (0.20)	8.52 (0.26)	6.46 (0.27)	7.38 (0.14)	7.09 (0.12)
W.Scale _{wt} ⁷	57.88 (0.04)	54.35 (0.07)	50.42 (0.07)	62.44 (0.02)	50.53 (0.08)

1 = Coefficient of variation. 2 = modal tuber size with respect to proportional tuber weight. 3 = modal tuber size with respect to proportional tuber weight. 4:5 = Weibull or scale shape of proportional tuber numbers. 6:7 = Weibull shape or scale parameter of proportional tuber weight.

of the TSD with respect to tuber numbers, ranged from 41 mm at Buttery Hill to 56 mm at Horse Foxhole and with respect to tuber weight, the scale was largest at Horse Foxhole (62.44 mm) and lowest at Buttery Hill (50.42 mm). This large variation between fields was not replicated within a field with the relative-form CV for the distribution scale ranging from 0.07 at Buttery Hill to 0.02 Horse Foxhole for both tuber size and weight. Within-field variation in TSD with respect to tuber number was higher when quantified as the shape of the distribution. The average shape of Weibull curves fitted on within-field TSD with respect to size ranged from 3.36 at Buttery Hill to 5.04 at Deaton 6. With respect to weight, higher shape values were observed, with the minimum at Buttery Hill (6.46) and maximum at HF7 (8.52). The higher shape values suggested that TSD with respect to tuber weight was more left-skewed than with respect to tuber number. Overall, more within-field variability was captured in the shape parameter than the scale parameter. The CV ranged from 0.12 at Deaton 6 and Horse Foxhole to 0.23 at Buttery Hill with respect to tuber number and up to 0.27 at

Buttery hill with respect to tuber weight. The CV of the scale parameter was consistently under 0.1 for both tuber number and weight, suggesting low variability in the modal tuber size or weight class despite large variability in the shape of the tail that influence the shape parameter.

Comparison of TSD functions

The distribution functions were fitted to the average proportional tuber weights at each location. The Gaussian and Gamma distributions predicted relatively lower probability densities at the modal tuber size compared to the two Weibull distributions (Fig. 2). The Weibull distributions tended to predict higher probability densities between 50 mm and 60 mm but the probability density quickly fell off towards the right tail, predicting low tuber yield in the oversized tuber size fractions (>65 mm). The Gaussian and Gamma distributions decrease towards the right tail and predict higher yields in the oversized fractions.




Fig. 2. The Gaussian, Gamma and Weibull distribution functions fitted to the average proportional tuber weights at HF7 (a), Buttery Hill (b), Crabtree Leasow (c), Deaton 6 (d) and Horse Foxhole (e).




Fig. 3. The Gaussian, Gamma and Weibull distribution functions fitted to the average proportional tuber numbers at HF7 (a), Buttery Hill (b), Crabtree Leasow (c), Deaton 6 (d) and Horse Foxhole (e).

The distribution functions were fitted to the proportional tuber numbers at all five locations (Fig. 3). The Gamma distribution tended to underestimate the TSD relative to the other three fitting methods. Apart from the right-skew at Buttery Hill, the TSD with respect to tuber number was roughly symmetric and the difference between the Gaussian and the two Weibull methods was not readily discernible visually. The Weibull shape at Crabtree Leasow, Buttery Hill, Horse Foxhole and Deaton 6 ranged from 4.7 to 5.0, suggesting left-skewed distribution (Table 3).

The log relative likelihood estimates of the overall distributions of the four tested models, relative to the overall distribution of the observed tuber numbers and tuber weights in each size fraction were calculated (Table 4). The confidence intervals of the estimates are also shown. With respect to tuber number, the Gaussian distribution was found to be the most plausible model at one out of the five locations (Crabtree Leasow) while the

Weibull distribution was the most plausible at Branston Booths, Buttery Hill and Deaton 6. The Weibull distribution with percentiles approach to parameter estimation was found to be most plausible at Horse Foxhole, where the tubers were sized in commercially practised main crop size fractions. With respect to tuber weight, the Weibull distribution with MLE was the most plausible model at Branston Booths, Buttery Hill, Crabtree Leasow and Deaton 6, with the highest relative likelihood to the observed discrete distribution as shown in Table 4. Similar to the distributions of the tuber numbers, the Weibull distribution with percentile-based parameter estimation was the most plausible model at Horse Foxhole with no overlapping confidence intervals at all five locations. Overall, the Gamma distribution was found to be the least plausible of the four tested models with respect to both tuber number and weight while the Gaussian distribution ranked second to the Weibull maximum

Table 4. Average log relative likelihood estimate and the margin of error (based on 95% confidence interval) of fitted Gaussian, Weibull, Gamma and Weibull Percentiles curves to potato TSDs at five, relative to the likelihood of the observed discrete distribution

Distr. ¹	HF7	B.Hill ²	C.Leasow ³	Deaton 6	H. Foxhole ⁴
Tuber Number					
Gaussian	-3.10 ± 0.03	-3.02 ± 0.02	-2.15 ± 0.03	-4.52 ± -0.03	-2.31 ± -0.01
W.MLE ⁵	-2.75 ± 0.03	-2.95 ± 0.02	-2.83 ± 0.02	-3.73 ± 0.02	-1.53 ± 0.02
Gamma	-5.27 ± 0.03	-3.47 ± 0.03	-2.74 ± 0.03	-8.71 ± 0.02	-2.48 ± 0.01
W.Perc ⁶	-3.68 ± 0.03	-3.86 ± 0.03	-2.42 ± 0.04	-5.38 ± 0.01	-0.03 ± 0.01
Tuber weight					
Gaussian	-345.31 ± 0.01	-272.14 ± 0.04	-160.71 ± 0.01	-249.75 ± 0.01	-3.87 ± 0.01
W.MLE	-197.33 ± 0.01	-196.74 ± 0.04	-110.32 ± 0.03	-222.35 ± 0.02	-2.70 ± 0.01
Gamma	-538.54 ± 0.01	-398.75 ± 0.01	-247.98 ± 0.01	-431.97 ± 0.02	-4.79 ± 0.03
W.Perc	-392.47 ± 0.01	-322.85 ± 0.03	-272.79 ± 0.02	-309.53 ± 0.02	-0.33 ± 0.01

1 = Distribution. 2 = Buttery Hill. 3 = Crabtree Leasow. 4 = Horse Foxhole. 5 = Weibull distribution with parameters estimated by MLE. 6 = Weibull distribution with parameters estimated by the percentiles approach.

likelihood estimates. The maximum likelihood estimates, particularly the Gamma and Weibull distributions, performed better than the Weibull percentiles approach in generalizing the predicted theoretical curve to the observed discrete distribution, on account of the fitting procedure which took full advantage of the discrete bins in the MLE. On the other hand, when less information (i.e. wider bins) was available as was the case at Horse Foxhole, the Weibull percentiles approach was more plausible than the maximum likelihood approach. While the overall generalization of the distribution to the observed discrete distribution is important, it is more crucial to maximize the log relative likelihood within the marketable tuber size fraction of the main crop (45–65 mm), where the mode of the distribution occurs. This is because the primary goal of modelling TSD is to estimate the tuber size fraction with the highest probability density (μ). This is used as an estimate of tuber size for the field. Following standard procedures from Travis (1987), the principles of scaling are adopted, such that the weight of an object is proportional to its volume and the volume is in turn proportional to the cubic power of its linear dimensions. Working backwards, the estimated

linear dimension (μ) can therefore be converted to a yield (weight) prediction, incorporating a pre-defined variety-specific tuber shape parameter and observed tuber number. The standard formula for this conversion is provided in Travis *et al.* (1987). The log relative likelihood of marketable tuber number and weight proportions were calculated (Table 5). With respect to tuber number, the results show that the Weibull percentiles approach generalizes better than all the three other approaches, on account of its high relative likelihood. The closed-form of the percentiles approach meant that the differences between the estimated proportions and the actual proportions were very small, with log relative likelihood approaching zero. For the maximum likelihood approaches, better generalization to the overall distribution as shown in Table 4 meant less goodness of fit in the marketable tuber portion where the mode of the distribution occurs. The same result was observed with respect to tuber weight, where the Weibull percentiles approach performed better than all three other approaches on account of its high log relative likelihood. With respect to both tuber number and weight, the Weibull distribution was also the most plausible of the three models fitted by

Table 5. Average log relative likelihood estimate and the margin of error (based on 95% confidence interval) of fitted Gaussian, Weibull, Gamma and Weibull Percentiles curves to the 45–65 mm size band of potato TSDs at five, relative to the likelihood of the observed discrete distribution

Distr. ¹	HF7	B.Hill ²	C.Leasow ³	Deaton 6	H. Foxhole ⁴
Marketable Tuber Numbers					
Gaussian	-4.05 ± 0.19	-2.10 ± 0.13	-1.45 ± 0.2	-6.81 ± 0.14	-13.97 ± 0.12
W.MLE ⁵	-2.64 ± 0.2	-2.23 ± 0.14	-0.79 ± 0.2	-4.50 ± 0.15	-11.49 ± 0.11
Gamma	-6.83 ± 0.17	-3.62 ± 0.12	-2.82 ± 0.18	-11.89 ± 0.13	-22.18 ± 0.1
W.Perc ⁶	0.00 ± 0.02	-0.17 ± -0.11	-0.12 ± -0.02	0.00 ± 0.02	0.00 ± 0.49
Marketable Tuber Weight					
Gaussian	-223.53 ± 0.03	-171.91 ± 0.03	-88.92 ± 0.03	-186.17 ± 0.02	0.53 ± 1.25
W.MLE	-29.73 ± 0.02	-66.03 ± 0.02	31.75 ± 0.03	-85.66 ± 0.02	0.53 ± 1.28
Gamma	-405.03 ± 0.02	-318.12 ± 0.02	-212.83 ± 0.02	-382.95 ± 0.02	0.53 ± 1.31
W.Perc	0.00 ± 0.02	-0.03 ± 0.01	-0.08 ± 0.02	0.00 ± 0.02	0.00 ± 0.05

1 = Distribution. 2 = Buttery Hill. 3 = Crabtree Leasow. 4 = Horse Foxhole. 5 = Weibull distribution with parameters estimated by MLE. 6 = Weibull distribution with parameters estimated by the percentiles approach.

Table 6. Root Mean Square Error (RMSE) of estimates from the Gaussian, Gamma and Weibull (Percentiles approach) benchmarked against the Weibull model with MLE

Location	gauRMSE ¹	gamRMSE ²	weiRMSE ³	shapeRMSE ⁴	scaleRMSE ⁵
Tuber number					
HF7	1.63	5.08	2.57	0.77	1.83
B.Hill ⁶	2.21	3.91	5.49	0.78	3.05
C. Leasow ⁷	1.55	4.36	2.53	1.21	1.82
Deaton 6	1.63	5.06	2.04	0.76	1.17
H.Foxhole ⁸	0.85	6.71	2.99	1.21	1.75
Tuber weight					
HF7	2.24	3.85	1.36	1.19	1.51
B.Hill	1.94	4.25	1.57	1.25	1.47
C. Leasow	2.17	3.84	1.35	1.07	1.70
Deaton 6	2.28	3.82	1.16	0.76	1.03
H.Foxhole	2.48	3.63	2.63	3.06	2.15

1 = RMSE of the mode of Gaussian model vs Weibull MLE, 2 = RMSE of the mode of the Gamma model vs Weibull MLE. 3 = RMSE of mode of the Weibull percentiles model vs Weibull MLE. 4 = RMSE of Weibull percentiles shape vs Weibull MLE. 5 = RMSE of Weibull percentiles scale vs Weibull MLE. 6 = Butterly Hill. 7 = Crabtree Leasow. 8 = Horse Foxhole.

MLE. This analysis showed that the Weibull distribution with the percentiles approach had the highest log relative likelihood of the observed marketable tuber profile, making it the best candidate for describing the TSD for the purposes of predicting the modal tuber size. However, the maximum likelihood estimate of the Weibull distribution proved to be a better model for generalizing the overall distribution outside the marketable component.

Using the Weibull distribution with MLE as a benchmark, the deviation of the other models from the benchmark in the prediction of the modal tuber size was as shown in Table 6. With respect to tuber weight, the Weibull percentiles approach yielded the lowest RMSE to the Weibull maximum likelihood approach's mode estimation at all locations except Horse Foxhole. However, as seen in Table 4, the percentiles approach was the most plausible model at Horse Foxhole, therefore its low RMSE approach was considered to be due to the lesser accuracy of the Weibull maximum likelihood approach at this location. These results were mirrored in the RMSEs of the modal tuber weight as shown in Table 6. Apart from Horse foxhole, the RMSE of the shape parameter between the maximum likelihood and percentiles approaches at all locations was within 1.5 units, showing that the percentiles approach approximated the shape of the distribution comparably to the benchmark. Similar observations were made for the scale parameter. At Horse Foxhole, higher RMSEs were observed, partially attributable to the lesser accuracy of the maximum likelihood approach. The analysis showed that the percentiles approach was comparable to the maximum likelihood approach in the estimation of the modal tuber size and yielded the highest log relative likelihood of the observed marketable yield. Therefore the distribution shape, scale and modal tuber size as estimated using the percentiles approach were tested for their predictability using within-field soil nutrient variation.

Linear modelling

The linear modelling results for the relationships between soil nutrients and TSD parameters with respect to tuber number are presented in Table 7. Stem density was also included as a fixed effect. For the shape parameter, no significant relationship was

observed with the stem density. This suggested that the proportional number of tubers falling into the pre-defined size bands was not affected by the stem density. However, P and N were observed to negatively associate with the shape parameter with statistical significance through the confidence intervals (Table 7). The strongest negative relationship with the shape parameter was observed with P, suggesting that P increased the proportional numbers of small-sized tubers. For the scale parameter, a significant negative relationship was observed with stem density, with a standardized regression coefficient of -0.98 , suggesting that increased stem densities led to a decrease in the overall range of the distribution. Positive relationships with the scale parameter were observed for P and N, suggesting that the two nutrients increased the overall range of the distribution, though only P was observed to hold a statistically significant relationship.


The overall effect of P on the TSD at the four sites was as shown in Fig. 4, where the average shape and scale parameter from low-P samples (P concentration less than the location mean) and high-P samples (P concentration higher than the location mean) were plotted to visualize the relationship. Overall, an increase in P at all four locations was associated with a shift of the TSD towards a right skew, leading to a lower probability density of the modal tuber size. As shown in Table 7, the mode of the distribution with respect to tuber number was also significantly related to the stem density and P, with a statistically significant regression coefficient of -2.37 for P. Setting the random effects to zero, the fixed-effects-only model had an nRMSE of 0.14 for the mode, suggesting that the expected modal tuber size across the field can be predicted to acceptable accuracy using stem density and soil nutrient information, with a Matern covariance structure fitted across the surface.

The regression coefficients of the shape, scale and mode variables of the TSD (with respect to tuber weight) as a function of stem density and the concentrations of soil nutrients were calculated (Table 8). The shape parameter was negatively related to stem density with a regression coefficient of -0.31 , showing that increasing stem numbers moderately associated with a shift of the TSD towards a right skew. The same association was also

Table 7. Linear modelling results for the relationships between soil nutrients (and stem density) and TSD parameters with respect to tuber number

Response	Predictors	Estimate	nRMSE _{fixed} ¹	delta AICc ²	D.F. ³	ICC1 ⁴
Shape ⁵	Intercept	4.35 ± 0.63	0.18	7.15	75.66	0.37
	Stem density	−0.04 ± 0.17				
	Phosphorus	−0.74 ± 0.33				
	Nitrogen	−0.29 ± 0.25				
Scale ⁵	Intercept	49.15 ± 7.22	0.13	14.77	72.19	0.87
	Stem density	−0.98 ± 0.63				
	Phosphorus	0.92 ± 0.27				
	Nitrogen	1.44 ± 1.16				
Mode ⁵	Intercept	45.71 ± 8.05	0.14	11.41	72.66	0.83
	Stem density	−1.08 ± 0.78				
	Phosphorus	−2.37 ± 1.99				
	Nitrogen	0.85 ± 1.40				

1 = Normalized Root Mean Square Error of the fixed effects model, with random effects set to zero. 2 = change in the conditional Akaike Information Criteria between the current model and the random intercept model. 3 = effective degrees of freedom. 4 = Intraclass correlation of the random effects. 5 = parameter determined from the best fitting model.

**Fig. 4.** Illustration of the effect of Phosphorus concentration on the TSD at Deaton 6 (a), Buttery Hill (b), HF7 (c), and Horse Foxhole (d).

observed with nutrients P and N, suggesting that increased soil nutrient concentrations in these soils were associated with an increase in the proportion of low-weight tubers. The relationships between the three variables and the shape parameter were all significant as evaluated through their non-zero confidence intervals.

The fixed-effects-only model fitted the data with an nRMSE of 0.22. Stem density also had a significant negative association with the scale parameter, which suggested that observations with relatively high stem densities had a smaller range of tuber sizes with a higher concentration of small tubers. Phosphorus

Table 8. Linear modelling results for the relationships between soil nutrients (and stem density) and TSD parameters with respect to tuber weight

Response	Predictors	Estimate	nRMSE _{fixed} ¹	delta AICc ²	D.F. ³	ICC1 ⁴
Shape ⁵	Intercept	7.46 ± 0.36	0.22	5.01	77	0.37
	Stem density	-0.31 ± 0.59				
	Phosphorus	-1.3 ± 0.62				
	Nitrogen	-0.89 ± 0.63				
Scale ⁵	Intercept	56.26 ± 5.79	0.09	12.69	74.23	0.72
	Stem density	-0.99 ± 0.78				
	Phosphorus	2.6 ± 1.83				
	Nitrogen	2.1 ± 1.17				
	Potassium	-0.75 ± 0.8				
Mode ⁵	Intercept	54.79 ± 6.28	0.21	11.94	75.14	0.79
	Stem density	-1.04 ± 0.69				
	Phosphorus	1.27 ± 1.57				
	Nitrogen	1.35 ± 1.07				

1 = Normalized Root Mean Square Error of the fixed effects model, with random effects set to zero. 2 = change in the conditional Akaike Information Criteria between the current model and the random intercept model. 3 = effective degrees of freedom. 4 = Intraclass correlation of the random effects. 5 = parameter determined from the best fitting model.

and N had significant positive relationships with the scale parameter, suggesting that the overall range of the distribution was increased with the increase of these soil nutrients. Potassium was also negatively related to the scale parameter. The fixed effects explained the variation in the scale parameter with an nRMSE of 0.09. The model fitted for the modal tuber size showed significant negative relationships between stem density and tuber size. Phosphorus and N both had positive relationships with the mode as shown by the positive regression coefficients in Table 8. However, the P relationship was not significant as evaluated the zero-coverage of the confidence interval. The fixed effect coefficients fitted the observed data with an nRMSE of 0.21.

Discussion

TSD model

The findings of the study showed that potato TSD was more consistent with the Weibull distribution than the Gamma and Gaussian distributions, consistent with Nemecek *et al.* (1996) and Bussan *et al.* (2007). The log relative likelihood analysis of the overall distribution showed that the maximum likelihood approaches generalized to the observed discrete proportions better than the percentiles approach and the Weibull distribution maximum likelihood approach was the best method for generalizing the shape and scale of the distribution. However, in the context of this study, the percentiles approach performed better at maximizing the likelihood of the tuber size fraction of interest (45–65 mm) because its closed-form offered the opportunity to maintain the likelihood of this desired size fraction accurately. Ultimately, the main goal of TSD modelling is to maximize the likelihood of modelling the correct modal tuber size of a collected sample, which will fall within a pre-defined tuber size interval. In the sample used in this study, the log relative likelihood analysis demonstrated comparable performance between the estimates for the marketable tuber size fraction determined using the Weibull percentiles approach and MLE. Additionally, the Weibull percentiles approach had the lowest RMSE for the prediction of the modal

tuber size, in comparison with the MLE approach, while maintaining the highest relative likelihood for the frequency of the marketable 45–65 mm size fraction. Unlike the percentiles approach, the accuracy of the MLE approach can be potentially improved with closer sample-grading intervals. However, closer grading entails more labour at grading time, which affects the adoption of the method in large-scale production environments, making the percentiles approach an attractive option. The relative likelihood for the marketable size range in the MLE method can also be improved by incorporating a weighted approach that provides more weight to the desired size fraction, a trading bias for precision where credible context-specific weight-selection methods can be produced (Hu and Zidek, 2002). Large variation in the Weibull shape parameter for TSD modelled against both tuber number and weight proportions shows the merit of adopting a flexible distribution function, in contrast to Nemecek *et al.* (1996) who suggested an average Weibull shape parameter of 2.3 as a general solution to simulate a right-skewed distribution. However, Nemecek *et al.* (1996) used tuber samples from potatoes grown for a seed market, which are desiccated early with less time allowed for tuber bulking. In the current study, a largely right-skewed distribution was observed at BATTERY Hill, which also had the smallest average tuber size. The Weibull distribution best modelled the BATTERY Hill crop as evidenced by the log relative likelihoods of both the overall distribution and the marketable portion. However, the shape parameter was 3.4 for TSD modelled after proportional tuber numbers and 6.46 for TSD modelled after proportional weight. This showed that proportional tuber numbers were roughly symmetrically distributed around the modal tuber size, but the proportional weights were slightly left-skewed because of a higher weight per tuber of the larger tubers.

Across all locations, the predicted modal tuber size with respect to tuber weight was larger than the modal tuber size with respect to tuber numbers. This is in line with principles of scaling whereby the weight of an object is proportional to its volume, which is in turn proportional to the cubic power of its linear dimensions, the principle behind the use of the modal tuber size for predicting yield in Travis *et al.* (1987). Therefore, in an

assumed Gaussian distribution of tuber numbers, the distribution of tuber weights can be expected to be slightly left-skewed with a higher modal size with respect to proportional tuber weights on account of an exponential increase of weight per tuber with increasing tuber size. These findings provide further evidence that the assumption of a Gaussian distribution as adopted by Travis (1987) does not universally hold true.

Struik *et al.* (1991) observed an initial right-skewed TSD with a modal tuber size gradually shifting to the right as more time was allowed for tuber bulking by shifting the harvest date. There is good evidence therefore that the overall shape of the distribution is influenced by the average tuber size, which depends on the time of observation, tuber bulking rate and the timing of harvest. It can be argued that the Gaussian (Sands and Regel, 1983), Log-normal (Marshall *et al.*, 1993) and Gamma (Aliche *et al.*, 2019) distributions, as well as the Weibull distribution with a fixed shape parameter, are all instance-specific realizations of an underlying dynamic distribution that is a function of the tuber bulking status at the time of observation. In turn, tuber bulking rates are controlled in part by the efficiency of source-sink transportation of photosynthates which depends on the plant nutrition status of the crop (Struik *et al.*, 1991). Allowing for a flexible shape parameter, the current results suggest that the Weibull distribution is the best estimator of the TSD curve at any stage of tuber development for the most representative prediction of the modal tuber size, which is the main purpose for modelling TSD.

A method for non-destructively modelling TSD is desirable for temporal monitoring of the changes in TSD. To achieve this, one of the parameters in the Weibull distribution needs to be modelled as a function of a non-destructively measurable variable. Both Nemecek *et al.* (1996) and Bussan *et al.* (2007) chose to model the scale parameter from empirical data, however, they did so for different reasons. Nemecek *et al.* (1996) suggested a fixed Weibull shape parameter, as they empirically observed that distribution fit was relatively invariant in their dataset. However, Bussan *et al.* (2007) explicitly modelled the scale parameter because it had a high correlation to measured stem and tuber density. In the current study, the shape parameter had higher within-field variability than the scale at all study sites as measured by the CV, regardless of the method by which TSD was measured (i.e. as a function of proportional tuber number or weight).

Effect of soil factors on the TSD

Literature on TSD is dominated by plant population studies, with the number of stems per unit area being reported to have a negative effect on indicators of TSD and tuber number (Knowles and Knowles, 2006; Bussan *et al.*, 2007; Aliche *et al.*, 2019). These past findings all support the hypothesis that higher stem numbers per unit area increase the tuber number per unit area at the expense of tuber size, leading to more uniform but smaller-sized tubers. The results of the multivariable regression are in general agreement with these findings as stem number per unit area had the most consistent negative regression coefficients with the modal tuber size and the scale parameter of TSD, regardless of whether TSD was modelled with respect to proportional tuber numbers or weight. A lower scale parameter associated with increased stem number means that there was a smaller probability density of large tubers, supporting the previous findings.

The standardized regression coefficients suggest for stem number suggest a large effect size of stem number on the modal tuber size and scale, to provide further support to the hypothesis that

stem density increase significantly affects tuber size. Stem density also had consistently negative relationships to the shape of the distribution but the effect size was low. This may be interpreted to mean that bulking rates of large tubers are reduced the presence of high stem numbers but the tuber filling is not altered to particularly favour any one of the other predefined size classes significantly.

The results suggest that the effects of soil macronutrients on TSD may be important, observing negative associations between P and the shape parameter with a high effect size for TSD measured with respect to tuber number or weight. This is consistent with findings that high P tests (25–33 mg/kg) may be associated with an increase in the proportion of small tubers in the TSD, attributed to increased vegetative growth at the expense of tuber bulking (see Birch *et al.*, 1967; Prummel and Barnau-Sijthoff, 1984; Sharma and Arora, 1987; Rosen and Bierman, 2008).

In the current study, the lowest concentration of available P was observed at Deaton 6 (41 mg/kg), with up to 100 mg/kg at Buttery Hill, which were much greater than the soil P tests at which negative effects were observed in the Rosen and Bierman (2008) study. While positive effects of P on tuber yield components have been reported from replicated experiments by Freeman *et al.* (1998), the responses were in low P soils and an asymptote was reached at 27 mg/kg for the Kennebec variety. The results give evidence of a significant negative relationship between P concentrations and the modal tuber size with respect to tuber numbers, meaning that the proportional number of smaller tubers increases as P concentration increases. However, the scale of the distribution with respect to both tuber number and weight was increased with an increase in P concentration. This meant suggests that P did not affect the tuber filling hierarchy but led to the induction of more tubers that remained within the small size fraction. The smaller tubers are expected to have a relatively low contribution to the overall yield therefore the positive effect of P on the scale of the distribution can be expected to positively affect the modal tuber size as observed in the study.

As a possible explanatory mechanism, potatoes are known to maintain P uptake even at high soil P tests like the ones observed in the current study (Jasim *et al.*, 2020), thereby accumulating inorganic P in the cytosol. Inorganic phosphate accumulation is inhibitory to the activity of ADP-glucose pyrophosphorylase and subsequently inhibit the starch synthesis and accumulation in sink organs (Crafts-Brandner, 1992; Kleczkowski, 1999; Tiessen *et al.*, 2002). A tuber-filling hierarchy has been previously shown whereby larger tubers grow the fastest and increase the spread of the TSD (Mackerron *et al.*, 1988), therefore high P concentrations can be expected to contribute to increased proportions of small tuber numbers without penalizing the scale of the distribution as shown in the study.

There have been a few previous studies where the effect of soil nutrients on TSD was systematically studied. Wurr *et al.* (1993) found that N had a significant effect of TSD measured as the spread (CV %) of tuber sizes (assuming a Gaussian distribution), however, they found no effect of P. The findings by Wurr *et al.* (1993) are also reflective of the design-intrinsic large concentration gradients of the nutrient treatments in controlled experiments, where N becomes a limiting nutrient hence its effects are emphasized. In the adequately fertilized sites of the current study where N was not a limiting factor for production, the evidence suggests that P equally contributes to the underlying model that explains within-field variation in TSD. Nitrogen was largely found to have the same effects on TSD parameters as P, with positive standardized

regression coefficients on the scale and mode with respect to tuber weight suggesting that N increases the variability in tuber size, in alignment with Wurr *et al.* (1993). High N was associated with a lower shape parameter, showing a particularly high effect size with respect to tuber weight. However, wide confidence intervals call for a more precise estimation of the relationship.

These findings mean that field variation in expected TSD can be constructed from soil data by modelling the shape parameter and scale parameter for the field. The scale and shape parameters can also be calibrated by mid-season yield digs. The beneficial properties of N and P on tuber yield are well documented in the literature and formulate the basis of crop fertilization modelling with the assumption of a Mitscherlich exponential growing process with a horizontal asymptote. The current findings suggest the existence of an inflection point after which over-fertilization results in a linear reduction in the shape of the TSD.

Conclusion

In conclusion, the study has shown that the shape parameter of the Weibull distribution determined using a linearized cumulative probability function provides an adequate index for describing the overall shape of TSD and performs better than the Gaussian and Gamma distributions in simulating observed TSD. The linearized formulae for the Weibull shape and scale presented in the current paper can be easily implemented in spreadsheet software at the farm level. Using the shape parameter, agronomists can improve the monitoring methods of the temporal shift in TSD from yield digs and (where large tubers are preferred) aim for symmetrical (shape ~ 3) or left-skewed (shape > 3) TSD. With the availability of Weibull cumulative probability functions in popular spreadsheet software, tuber numbers in any discrete size grades can be calculated from the modelled shape and scale parameter. From the current study, the shape parameter had larger within-field variability than the scale parameter and was significantly affected by excess P and N. Ultimately, high-intensity soil maps of these elements can enable high-resolution modelling of spatial TSD variation. While requirements for high-intensity soil sampling remain prohibitive, modelling of soil variability using co-kriging proxies like apparent electrical conductivity becomes important and relevant for generating high-resolution field variation for decision-support. Subsequent studies to validate these findings in more environments are recommended, as well as controlled studies to investigate the general point of inflection at which additional fertilization becomes detrimental to TSD.

Acknowledgements. Acknowledgement to the Crops and Environment Research Centre of Harper Adams University and Branston Limited for providing the experimental sites for this study. Many thanks to William Johnson, Andrew Beacham, Kevin Jones, Gary Weston, Grace Milburn and Joseph Crosby for their technical help.

Financial support. This work was supported by the Agriculture and Horticulture Development Board (Stoneleigh Park, Kenilworth, Warwickshire, CV8 2TL, United Kingdom) [Project Number 11140054].

Conflicts of interest. The authors declare there are no conflicts of interest.

Ethical standards. Not applicable.

References

- Aliche EB, Oortwijn M, Theeuwens TPJM, Bachem CWB, van Eck HJ, Visser RGF and van der Linden CG (2019) Genetic mapping of tuber size distribution and marketable tuber yield under drought stress in potatoes. *Euphytica* **215**, 1–19.
- Allison MF, Fowler JH and Allen EJ (2001) Responses of potato (*Solanum tuberosum*) to potassium fertilizers. *Journal of Agricultural Science* **136**, 407–426.
- Andrade MHML, da Silva ALBR, Pesantes LG, Christensen CT and Zotarelli L (2021) Seed piece spacing for early-maturing table-stock potato grown under subtropical conditions. *American Journal of Potato Research* **98**, 246–254.
- Arsenault WJ, LeBlanc DA, Tai GCC and Boswall P (2001) Effects of nitrogen application and seedpiece spacing on yield and tuber size distribution in eight potato cultivars. *American Journal of Potato Research* **78**, 301–309.
- Birch JA, Devine JR, Holmes MRJ and Whitear JD (1967) Field experiments on the fertilizer requirements of maincrop potatoes. *The Journal of Agricultural Science* **69**, 13–24.
- Buhrig W, Thornton MK, Olsen N, Morishita D and McIntosh C (2015) The influence of ethephon application timing and rate on plant growth, yield, tuber size distribution and skin color of red LaSoda potatoes. *American Journal of Potato Research* **92**, 100–108.
- Bussan AJ, Mitchell PD, Copas ME and Drilias MJ (2007) Evaluation of the effect of density on potato yield and tuber size distribution. *Crop Science* **47**, 2462–2472.
- Caldiz DO, Viani PG, Giletto CM, Zamuner EC and Echeverría HE (2018) Improving yield and quality of processing potato crops grown in the Argentinian pampas: the role of N, P and S and their impact on CO₂ emissions. *Potato Research* **61**, 147–168.
- Champely S, Ekstrom C, Dalgaard P, Gill J, Weibelzahl S, Anandkumar A, Ford C, Volcic R and De Rosario H (2018) Package “pwr”: basic functions for power analysis. CRAN Repository, Version: 1.3-0, Software: <https://cran.r-project.org/web/packages/pwr/>
- Cohen J (1988) *Statistical Power Analysis for the Behavioural Sciences*. Hillsdale, NJ: Lawrence Earlbaum Associates.
- Costa JFF, Giasson É, da Silva EB, Coblinski JA and Tiecher T (2020) Use of color parameters in the grouping of soil samples produces more accurate predictions of soil texture and soil organic carbon. *Computers and Electronics in Agriculture* **177**, 105710.
- Crafts-Brandner SJ (1992) Phosphorus nutrition influence on starch and sucrose accumulation, and activities of ADP-glucose pyrophosphorylase and sucrose-phosphate synthase during the grain filling period in soybean. *Plant Physiology* **98**, 1133–1138.
- Delignette-Muller ML and Dutang C (2015) Fitdistrplus: an R package for fitting distributions. *Journal of Statistical Software* **64**, 1–34.
- Dickins JC, Harrap FEG and Holmes MRJ (1962) Field experiments comparing the effects of muriate and sulphate of potash on potato yield and quality. *The Journal of Agricultural Science* **59**, 319–326.
- Dubey SD (1967) Some percentile estimators for Weibull parameters. *Technometrics* **9**, 119–129.
- ESRI (2018). ArcGIS Desktop. ArcGIS Blog.
- Freeman KL, Franz PR and de Jong RW (1998) Effect of phosphorus on the yield, quality and petiolar phosphorus concentrations of potatoes (cvv. Russet Burbank and Kennebec) grown in the krasnozem and duplex soils of Victoria. *Australian Journal of Experimental Agriculture* **38**, 83–93.
- Gao X, Shaw WS, Tenuta M and Gibson D (2018) Yield and nitrogen use of irrigated processing potato in response to placement, timing and source of nitrogen fertilizer in Manitoba. *American Journal of Potato Research* **95**, 513–525.
- Henderson R (1965) Effects of type of potash and rates of nitrogen, phosphate and potassium in fertilisers on growth, yield and quality of potato crops. *Journal of the Science of Food and Agriculture* **16**, 480–488.
- Hu F and Zidek JV (2002) The weighted likelihood. *Canadian Journal of Statistics* **30**, 347–371.
- Jackson E, Farrington DS and Henderson K (1986) *The Analysis of Agricultural Materials: A Manual of the Analytical Methods Used by the Agricultural Development and Advisory Service*. London: Reference Book, ADAS, Ministry of Agriculture, Fisheries and Food.
- Jasim A, Sharma LK, Zaen A, Bali SK, Buzza A and Alyokhin A (2020) Potato phosphorus response in soils with high value of phosphorus. *Agriculture* **10**, 264.

- Keijbets MJH** (2008) Potato processing for the consumer: developments and future challenges. *Potato Research* **51**, 271–281.
- Kirsten WJ and Hesselius GU** (1983) Rapid, automatic, high-capacity dumas determination of nitrogen. *Microchemical Journal* **28**, 529–547.
- Kleczkowski LA** (1999) A phosphoglycerate to inorganic phosphate ratio is the major factor in controlling starch levels in chloroplasts via ADP-glucose pyrophosphorylase regulation. *FEBS Letters* **448**, 153–156.
- Knowles NR and Knowles LO** (2006) Manipulating stem number, tuber set, and yield relationships for northern- and southern-grown potato seed lots. *Crop Science* **46**, 284–296.
- Lai C-D, Murthy DN and Xie M** (2006) Weibull distributions and their applications. In Pham Hoang (ed.), *Springer Handbook of Engineering Statistics*. London: Springer, pp. 63–78.
- Lindsey JK** (1974) Comparison of probability distributions. *Journal of the Royal Statistical Society: Series B (Methodological)* **36**, 38–47.
- Love SL and Thompson-Johns A** (1999) Seed piece spacing influences yield, tuber size distribution, stem and tuber density, and net returns of three processing potato cultivars. *HortScience* **34**, 629–633.
- Ly A, Marsman M, Verhagen J, Grasman RPPP and Wagenmakers E-J** (2017) A tutorial on fisher information. *Journal of Mathematical Psychology* **80**, 40–55.
- Machakaire ATB, Steyn JM, Caldiz DO and Haverkort AJ** (2016) Forecasting yield and tuber size of processing potatoes in South Africa using the LINTUL-potato-DSS model. *Potato Research* **59**, 195–206.
- Mackerron DKL, Marshall B and Jefferies RA** (1988) The distributions of tuber sizes in droughted and irrigated crops of potato. II. Relation between size and weight of tubers and the variability of tuber-size distributions. *Potato Research* **31**, 279–288.
- Marshall B, Holwerda HT and Struik PC** (1993) Synchronisation of tuber growth in potato (*Solanum tuberosum*): a statistical model. *Field Crops Research* **32**, 343–357.
- Minasny B and McBratney AB** (2005) The Matérn function as a general model for soil variograms. *Geoderma* **128**, 192–207.
- Mponela P, Snapp S, Villamor GB, Tamene L, Le QB and Borgemeister C** (2020) Digital soil mapping of nitrogen, phosphorus, potassium, organic carbon and their crop response thresholds in smallholder managed escarpments of Malawi. *Applied Geography* **124**, 102299.
- Nemecek T, Derron JO, Roth O and Fischlin A** (1996) Adaptation of a crop-growth model and its extension by a tuber size function for use in a seed potato forecasting system. *Agricultural Systems* **52**, 419–437.
- Page AL** (1982) *Methods of Soil Analysis-Part 2: Chemical and Microbiological Properties*, 2nd Edn. Madison, USA: Am. Soc. Agron. Inc. Publ.
- Porter GA and Sisson JA** (1991) Petiole nitrate content of Maine-grown Russet Burbank and Shepody potatoes in response to varying nitrogen rate. *American Potato Journal* **68**, 493–505.
- Prummel J and Barnau-Sijthoff P** (1984) Optimum phosphate and potassium levels in potato tops. *Fertilizer Research* **5**, 203–211.
- R Core Team** (2019). R: A language and environment for statistical computing. R Foundation for Statistical Computing. Vienna, Austria.
- Rietra RPJJ, Heinen M, Dimkpa CO and Bindraban PS** (2017) Effects of nutrient antagonism and synergism on yield and fertilizer use efficiency. *Communications in Soil Science and Plant Analysis* **48**, 1895–1920.
- Rosen CJ and Bierman PM** (2008) Potato yield and tuber set as affected by phosphorus fertilization. *American Journal of Potato Research* **85**, 110–120.
- Rousset F and Ferdy JB** (2014) Testing environmental and genetic effects in the presence of spatial autocorrelation. *Ecography* **37**, 781–790.
- Sands PJ and Regel PA** (1983) V. A simple model for predicting graded yields. *Field Crops Research* **6**, 25–40.
- Schippers PA** (1968) The influence of rates of nitrogen and potassium application on the yield and specific gravity of four potato varieties. *European Potato Journal* **11**, 23–33.
- Sharma UC and Arora BR** (1987) Effect of nitrogen, phosphorus and potassium application on yield of potato tubers (*Solanum tuberosum* L.). *The Journal of Agricultural Science* **108**, 321–329.
- Simmons KE and Kelling KA** (1987) Potato responses to calcium application on several soil types. *American Potato Journal* **64**, 119–136.
- Stein ML** (1999) *Interpolation of Spatial Data*. New York: Springer.
- Struik PC, Haverkort AJ, Vreugdenhil D, Bus CB and Dankert R** (1990) Manipulation of tuber-size distribution of a potato crop. *Potato Research* **33**, 417–432.
- Struik PC, Vreugdenhil D, Haverkort AJ, Bus CB and Dankert R** (1991) Possible mechanisms of size hierarchy among tubers on one stem of a potato (*Solanum tuberosum* L.) plant. *Potato Research* **34**, 187–203.
- Taylor JA, Chen H, Smallwood M and Marshall B** (2018) Investigations into the opportunity for spatial management of the quality and quantity of production in UK potato systems. *Field Crops Research* **229**, 95–102.
- Tiessen A, Hendriks JHM, Stitt M, Branscheid A, Gibon Y, Farré EM and Geigenberger P** (2002) Starch synthesis in potato tubers is regulated by post-translational redox modification of ADP-glucose pyrophosphorylase. *The Plant Cell* **14**, 2191–2213.
- Travis KZ** (1987) Use of a simple model to study factors affecting the size distribution of tubers in potato crops. *The Journal of Agricultural Science* **109**, 563–571.
- van Dijk LCM, Lommen WJM, de Vries ME, Kacheyo OC and Struik PC** (2021) Hilling of transplanted seedlings from novel hybrid true potato seeds does not enhance tuber yield but can affect tuber size distribution. *Potato Research* **64**, 353–374.
- Westermann DT and Kleinkopf GE** (1985) Phosphorus relationships in potato plants. *Agronomy Journal* **77**, 490–494.
- Wurr DCE, Fellows JR, Lynn JR and Allen EJ** (1993) The impact of some agronomic factors on the variability of potato tuber size distribution. *Potato Research* **36**, 237–245.
- Yang Y, Luo Y, Lu M, Schädel C and Han W** (2011) Terrestrial C:N stoichiometry in response to elevated CO₂ and N addition: a synthesis of two meta-analyses. *Plant and Soil* **343**, 393–400.