

Lines in the landscape

by Clifford, C., Bieroza, M., Clarke, S.J., Pickard, A., Stratigos, M.J., Hill, M.J., Raheem, N., Tatariw, C., Wood, P.J., Arismendi, I., Audet, J., Aviles, D., Bergman, J.N., Brown, A.G., Burns, R.E., Connolly, J., Cook, S., Crabot, J., Cross, W.F., Dean, J.F., Evans, C.D., Fenton, O., Friday, L., Gething, K.J., Giannico, G., Habib, W., Maher Hasselquist, E., Heili, N.M., van der Knaap, J., Kosten, S., Law, A., van der Lee, G.H., Mathers, K.L., Morgan, J.E., Rahimi, H., Sayer, C.D., Schepers, M., Shaw, R.F., Smiley Jr, P.C., Speir, S.L., Strock, J.S., Struik, Q., Tank, J.L., Wang, H., Webb, J.R., Webster, A.J., Yan, Z., Zivec, P. and Peacock, M.

Copyright, publisher and additional information: Publishers' version distributed under the terms of the [Creative Commons Attribution License](#)

[DOI link to the version of record on the publisher's site](#)



**Harper Adams
University**

Clifford, C., Bieroza, M., Clarke, S.J., Pickard, A., Stratigos, M.J., Hill, M.J., Raheem, N., Tatariw, C., Wood, P.J., Arismendi, I., Audet, J., Aviles, D., Bergman, J.N., Brown, A.G., Burns, R.E., Connolly, J., Cook, S., Crabot, J., Cross, W.F., Dean, J.F., Evans, C.D., Fenton, O., Friday, L., Gething, K.J., Giannico, G., Habib, W., Maher Hasselquist, E., Heili, N.M., van der Knaap, J., Kosten, S., Law, A., van der Lee, G.H., Mathers, K.L., Morgan, J.E., Rahimi, H., Sayer, C.D., Schepers, M., Shaw, R.F., Smiley Jr, P.C., Speir, S.L., Strock, J.S., Struik, Q., Tank, J.L., Wang, H., Webb, J.R., Webster, A.J., Yan, Z., Zivec, P. and Peacock, M. (2025) 'Lines in the landscape' *Communications earth & environment*, 6, Article number 693.

<https://doi.org/10.1038/s43247-025-02699-y>

Lines in the landscape

Check for updates

Chelsea Clifford ^{1,44}✉, Magdalena Bieroza ², Stewart J. Clarke ³, Amy Pickard⁴, Michael J. Stratigos ⁵, Matthew J. Hill⁶, Nejem Raheem⁷, Corianne Tatariw ⁸, Paul J. Wood⁹, Ivan Arismendi ¹⁰, Joachim Audet¹¹, Daniel Aviles ¹², Jordanna N. Bergman¹³, Anthony G. Brown^{14,15}, Rachel Eleanor Burns¹⁶, John Connolly¹⁷, Sarah Cook¹⁸, Julie Crabot ¹⁹, Wyatt F. Cross²⁰, Joshua F. Dean ²¹, Chris D. Evans ²², Owen Fenton²³, Laurie Friday²⁴, Kieran J. Gething ²⁵, Guillermo Giannico¹⁰, Wahaj Habib ¹⁷, Eliza Maher Hasselquist²⁶, Nathaniel M. Heili²⁰, Judith van der Knaap ²⁷, Sarian Kosten²⁷, Alan Law²⁸, Gea H. van der Lee²⁹, Kate L. Mathers⁹, John E. Morgan ²¹, Hamidreza Rahimi²⁴, Carl D. Sayer³⁰, Mans Schepers³¹, Rosalind F. Shaw³², Peter C. Smiley Jr.³³, Shannon L. Speir³⁴, Jeffrey S. Strock^{35,36}, Quinten Struik²⁷, Jennifer L. Tank³⁷, Hao Wang ³⁵, Jackie R. Webb ^{38,39}, Alex J. Webster⁴⁰, Zhifeng Yan ⁴¹, Peta Zivec ²⁶ & Mike Peacock ^{42,43,44}✉

Ditches (linear constructions which store and/or move water where humans prefer it to go), via irrigation, drainage, and power, have helped drive the development of human societies. Now, ditches and other linear channels, typically carrying water, are numerous and found on every continent. Their form varies widely with use, which includes land drainage, irrigation, transportation, and boundary marking. Ditches support and shape biogeochemical cycles, biotic communities, and human societies, at multiple spatiotemporal scales. However, ditches are frequently overlooked by researchers in many disciplines. Here, we review the largely unrecognized role that ditches play in environmental processes and human societies. The effects of ditches can be both positive (e.g., biodiversity refuges, water for food production, nutrient retention) and negative (e.g., greenhouse gas emissions, dispersal of pollutants). We call for future management to consider and enhance the multifunctional role that ditches can deliver at the landscape-scale.

Human societies, globally, have enacted control of hydrological systems through drainage and irrigation, with evidence of linear waterways dating as far back as 8000 years¹. Ditches enabled the expansion of agriculture and settlements, by both supplying and excluding water, and allowed early humans to produce a surplus of food, which allowed part of society to be freed from agricultural labor. Thus, ditches provided the opportunities for societies to focus time and resources on cultural and commercial endeavors². Today, ~40% of global food production relies on irrigation and ~15% on drainage, and these dependencies are expected to increase³. Although ditches are widely distributed and found on every continent (even Antarctica⁴), their global extent is poorly quantified, with drained and irrigated cropland estimates varying between 130–200 Mha and 270–300 Mha^{3,5}. In the Northern Hemisphere, forestry drainage occupies an additional 15 Mha⁶ (Fig. 1C). The extent of ditch networks at the national scale can be extremely large (e.g., 300,000 km in the Netherlands⁷; 800,000 km in Sweden⁸) and a rough estimate of the global surface area of drainage ditches alone is 1.4–10.7 Mha⁹.

As ditch and irrigation networks have grown globally, so have their effects on natural hydrological cycles; they have drained and degraded wetlands¹⁰ and altered water flows through ecosystems at the landscape scale and beyond¹¹. At the same time, the existence of non-human life has also become increasingly interconnected with these channels^{12–14}. Ditches are palimpsests, echoes of past landscapes overwritten by human actions and naturalized again, embodying both creation and destruction. They challenge the human-nature dichotomy¹⁵, both of, and not of, natural waters. Simultaneously, ditches constitute an integral part of the hydrological network supporting remnants of aquatic ecosystems¹⁶ and exemplifying novel ecosystems¹⁷. Ditches often mimic natural waterways, whether by design or inadvertently, and yet sometimes surprise scientists by behaving differently⁹. In turn, they have the potential to serve as models for experimentation, which can provide insights into how natural ecosystems may respond to global change^{18,19}. Ditch networks can vary widely in their characteristics across space and time, both among and within individual channels²⁰, which can switch between terrestrial and aquatic states. This range, coupled with a

A full list of affiliations appears at the end of the paper. ✉e-mail: ccclifford@vims.edu; chelseaclifford@gmail.com; m.peacock@liverpool.ac.uk; michael.peacock@slu.se




Fig. 1 | Ditches vary in form and function. **A** an urban drainage ditch in Venice, Florida, USA. **B** the Oudegracht (“old canal”, dating from 1100 s) in urban Utrecht, the Netherlands, **C** an intermittent ditch in forestry, Sweden, **D** a ditch in a drained peatland converted to grassland, Iceland, **E** a roadside drainage in Ames, Iowa, USA,

F a drainage ditch in intensively-managed arable peatland, UK, **G** monastic (latrine) drain at the medieval Norton Priory, UK, **H** water management canal in an Acacia plantation, Indonesia. Photo credits: Chelsea Clifford (**A**, **E**), Mike Peacock (**B**, **F**, **G**), Eliza Maher Hasselquist (**C**), Chris Evans (**D**, **H**).

high degree of human control, makes ditches ideal systems for adaptively responding to global environmental change and to shifting human needs and wants¹⁸. Yet the diverse ecosystem services from ditches often conflict when they sometimes could synergize to deliver multifunctionality (i.e., multiple ecosystem services simultaneously)^{20,21}. Maximizing positive synergies requires collaboration between disciplines and stakeholders to optimize ditch potential and avoid pitfalls¹⁸. Unfortunately, despite their fundamental importance to society and potential to supplement ecosystem services, ditches remain understudied and undervalued^{21,22}.

For the purposes of this paper, we define “ditch” as a narrow linear channel on Earth’s surface constructed to store and/or convey water where humans prefer it to go. However, the vague boundaries of this definition are somewhat arbitrary; defining ditches is surprisingly non-trivial. Typically, they are narrow, but some can be wider than 25 m²³, and individual channels may run for just a few metres or for hundreds of kilometres in length. Although “ditch” is an English word, other languages classify what English-speakers might call *ditches* differently (see Table 1 for collated definitions). Therefore, even though ditches exist globally, the word “ditch” may not translate well across cultures. Even the English language might use different words for ditches depending on their dominant functions including: irrigation and drainage (e.g., rhyne, gripe, catchwater, gutter, dyke, conduit), water storage, power (e.g., leat), burial, bioreaction, transport (e.g., canal, waterway), defense (e.g., moat), livestock control, and boundary-marking, including as barriers (e.g., ha-ha) to encroachment. The purposes of ditches, the way they have been constructed and managed, and their resultant characteristics and classification vary over time and space (Fig. 1, Supplementary Table 1). Furthermore, the perception and value of ditches may influence how we classify them. The word “ditch” itself has a negative connotation within the English language both as the verb to ‘ditch’, meaning get rid of, and as exemplified by idioms like “ditching” something unwanted, and “dull as ditchwater.” This perspective may lead humans to avoid defining features we value as ditches, reaching instead for more positively associated terms such as “canal,” “stream,” or “blue infrastructure.” From an environmental perspective, ditches’ inherent artificiality can devalue them^{13,24}. That said, even artifice can be murky non-dichotomous, as in the case of

channelized streams. Naturalization away from constructed form often occurs over time, especially in the absence of repeated dredging¹³, and ultimately ditches can become archaeological features²⁵. Similarly, ditches and their subtypes fall somewhere within gradients in many of their physical characteristics, including size, network position, composition of bed material, connectivity to other waterbodies and floodplains, flow direction and speed, and even ability to hold water. The precise boundary around all ditch-like systems within these gradients and categories is, as we found, nearly impossible to agree upon, and ultimately arbitrary. That observation itself points towards ditches’ actual status as indivisible component of socio-ecosystems (an ecosystem and its associated human actors). So, for the purposes of this article, we shall not endeavor to precisely corral “ditches” from “not ditches.” We acknowledge that even our working definition will vary somewhat over the course of this article, in part as it crosses disciplinary lines and corresponding definitions.

In this paper, we present a multi-disciplinary perspective on how and why ditches are important. We review the state of knowledge pertaining to the different aspects of ditches including physical, biotic (excluding humans), chemical, and human dimensions. We also recommend future research and management considerations and outline knowledge gaps. Many of these future considerations presented will be opportunities, for ditches are rife with these; the potential for ditches to be adaptively managed for greater diversity of ecosystem services^{13,18,21}. Given that people will continue to construct and maintain ditches, we as researchers have an opportunity to positively influence their design, use, and management²⁶. At the same time, we can gain more broadly applicable knowledge from these distinctive, but common socio-ecosystems¹⁹. If we only let them, ditches can serve as a multi-tool for both humans and other species to survive and thrive under global environmental change¹⁸, helping to achieve sustainable development goals²⁷, and addressing a myriad future challenges. We hereby invite researchers who have not yet done so to meet the multitudes of the ditch.

Human dimension

Our working definition of ditches necessitates human conception, execution, and embodiment. In other words, without people playing

Table 1 | Summary of ditch definitions collated from published work

Type	Definition	Reference
Agricultural	"Man-made channels created primarily for agricultural purposes; and which usually: (i) have a linear platform, (ii) follow linear field boundaries, often turning at right angles, and (iii) show little relationship with natural landscape contours."	Williams et al. ²³⁷ , Brown et al. ²³⁸ , Davies et al. ²³⁹ , Davies et al. ²⁴⁰ , Clarke ²⁰³ , Shaw et al. ²⁴¹ , Hill et al. ¹⁸⁷ , Biggs et al. ¹⁶ , Bubikova and Hrivnak ²⁴² , Nakano and Morii ²⁴³ , Williams et al. ²⁴⁴
Agricultural	"Ditches are defined as artificial, linear channels < 3 m wide which follow anthropogenic boundaries (e.g., field margins). Drains are larger features (> 5 m wide) which display otherwise similar characteristics."	Gething and Little ²⁴⁵
Agricultural	"Dutch ditches are linear water bodies typically several metres wide and up to 1 m deep."	Verdonschot et al. ⁷
Agricultural	"Drainage ditches are small, stagnant, line-shaped water bodies, dug to improve rainwater run off and regulate the groundwater level of surrounding agricultural areas."	Verdonschot et al. ²⁴⁶
Agricultural	"Ditches are linear elements with a high edge ratio that are subjected to an intensive exchange of matter and organisms from the surrounding terrestrial matrix. Most of the ditches are likely to be relatively shallow with marked fluctuations in water levels and a higher probability of drying out during summer. Finally, ditches are regularly managed for efficient drainage."	Herzon and Helenius ²¹
Agricultural	"Drainage ditches are limited to those structures created to drain production acreage."	Cooper et al. ²⁴⁷
Agricultural	"Farm ditches are human-made linear elements that constitute the upstream parts of the permanent hydrographic networks in agricultural landscapes. Primarily implanted within farmed landscape to collect surface and subsurface water in order to drain excess water and/or to prevent soil erosion..."	Dollinger et al. ²⁰
Agricultural	"Agricultural drainage ditches are essentially headwater streams, which, like capillaries, act as direct links between agricultural fields and naturally occurring streams and rivers."	Fu et al. ²⁴⁸
Agricultural	"Ditches were defined as open field drains which flow into streams and are generally unmapped."	Shore et al. ²⁴⁹
Agricultural	"'Ditch' is used to describe systems either created or maintained by human activities in order to increase water conveyance; whereas 'drainage' refers to the practice water removal, or, when used in conjunction with 'network' or 'system,' describes the entirety of streams and ditches modified for water conveyance."	Pierce et al. ²⁵⁰
Agricultural	"Ditches... artificial linear water bodies whose depth and flow are regulated by sluice gates and pumping stations for the purposes of water-level management."	Watson & Ormerod ²⁰⁸
Agricultural	"Artificial channels built for agriculture irrigation purposes, generally have a regular U-shape and are approximately 0.5–3 m wide and 0.5–1 m deep distributed around farmland in agricultural regions."	Sun et al. ¹⁹⁵
Forest	"Headwater streams were classified as ditches if they were perfectly straight, if they made unnaturally sharp turns (e.g., 90° turns), or if they were clearly part of ditch networks (i.e., numerous parallel watercourses, geometric drainage networks)."	Peacock et al. ¹⁶⁵
Roadside	"Grassed roadside drainage ditches are shallow, open vegetated channels that are designed to convey stormwater runoff to storm sewers or receiving water bodies."	Ahmed et al. ²⁵¹
General	"Drainage ditches are small, linear water bodies, usually <1.5 m deep and several metres wide, situated both in lowland and in highland zones."	Nsenga Kumwimba et al. ²⁵²
General	"A long, narrow excavation artificially dug in the ground; especially an open and usually unpaved waterway, channel, or trench for conveying water for drainage or irrigation, and usually smaller than a canal. Some ditches may be natural watercourses."	European Protection Agency ²⁵³
General	"Ditches, irrigation channels and water supply canals, are constructed linear waterways, although their physical characteristics and function may vary widely."	Peacock et al. ⁹

Note: this is not intended to be fully comprehensive. In the text, we mention the problems that arise when considering the English word "ditch" and other languages. For example, in Dutch, "ditch" best translates as "sloot". A sloot is defined as an artificial permanent linear water body, exhibiting a maximum width of 8 meters and usually not more than 1.5 m deep with negligible flow (<5 cm/s). Meanwhile, a greppel, is usually smaller than a sloot, with only artificial and intermittent or ephemeral flow, but would also translate to English as "ditch." We lump the two terms, sloot and greppel, along with several other independent Dutch words like (urban) wadi and goot, together as ditches in English.

a vital role at some point in their life history, a ditch is not a ditch. This fact may in great part be responsible for the lack of attention ditches have received in the natural sciences—they are not 'natural' when natural is understood as lacking human influence. In contrast, ditches have been a major topic of historical and archaeological inquiry (Fig. 1G), which traces ditches through time for at least the

past 8000 years in different places around the globe precisely because they are clear signs of human activity. A range of social sciences have also addressed governance processes, how ditches affect human well-being, and a suite of ditch-related legal issues^{28,29}. Ditches are thus prime examples of socio-ecosystems; complex and integrated systems that include both the ecological and sociological environments, and

the interactions between them³⁰. Given this, the human part of the equation is essential to understand the whole.

Within our broad definition of ditches, irrigation (i.e., applying water to land for agriculture) is well-established in archaeological and historical scholarship and has been implicated as a key driver of state formation (i.e., the creation of states) and economic/social stratification in early agricultural societies^{31,32}. Evidence for ditch-based irrigation systems are found in Mesopotamia from as early as the 6th millennium BC^{33–36}. Early and indigenous ditch-based irrigation systems have also been examined across Asia^{37,38} and North^{29,39,40}, Central⁴¹ and South America⁴². Irrigation ditches of the past few centuries are somewhat less prominently studied archaeologically than older ones (but see De Meulemeester⁴³). However, they are the subject of more substantial documentary research⁴⁴. It should be noted that ditches and other hydrological controls are not exclusive to agricultural societies; fisher-hunter-gather societies have also constructed ditches to facilitate fishing⁴⁵. Ditches are often complex archaeological features due to burial and preservation processes related to long-use histories, especially in prehistoric contexts⁴⁶. There are also strong scholarly traditions in the theoretical underpinning of archaeological and historical interpretation of irrigation ditches, sometimes borrowing ecological concepts, for example viewing ditches through the lens of niche construction theory⁴⁷ or coevolution⁴⁸. Canals (Fig. 1B) and leats (ditches which specifically serve mills as either in- or outflows) have also been studied as part of power, transport, and urban histories⁴⁹. Indeed, in Europe mill streams and their associated leats and sluices can be considered an example of coevolutionary socio-ecohydrological systems (*sensu* Sivapalan & Blöschl⁴⁸), whereby over hundreds of years natural streams were modified to drive mills, which were later closed, leaving behind a hydrological legacy of remnant constructed channels⁵⁰.

Globally, 21% of wetlands have been anthropogenically drained¹⁰, largely via drainage ditches (i.e., ditches that remove excess water from the land) (Fig. 1D, F, H), resulting in a massive loss of natural wetlands. Despite this, drainage ditches have been less often archaeologically or historically studied than irrigation ditches, although this is changing with the global rise in the importance and value placed on peatland landscapes for biodiversity and carbon sequestration⁵¹. While historical attention to drainage is common⁵², research usually occurs at broader scales than examining or mapping the ditches themselves, with a view to understanding wider society, institutions, and worldviews. For example, the 17th century drainage of the Fens in eastern England has been the subject of extensive research, examining its economic impact⁵³, the politics of wetland drainage⁵⁴, and its public health implications⁵⁵. Taking drainage ditches' relational and land parcel-form into account also makes it possible to identify former hydrological conditions. For example, drainage ditches in now fully leveled reclaimed bog polder in the Netherlands sometimes still show a wedge-shaped structure, revealing their original orientation towards the top of the now missing domes of raised bogs⁵⁶ (Fig. 2). The drainage of wetlands, particularly over the last 1000 years for agriculture created both ditches and hedges which preserved relicts of past woodland cover and now form key havens of biodiversity in such environments^{57,58}.

Drainage ditches are usually understood archaeologically as threatening wetland archaeology through desiccation and other changes to burial environments⁵⁹. This threat has prompted numerous regional assessments of drainage ditch-driven wetland loss as a way to estimate where and what wetland archaeology remains well preserved in-situ⁶⁰. The archaeological and palaeoecological value of wetlands and ditches (a cultural ecosystem service in its own right) often comes primarily through anoxic conditions allowing for the survival of materials that in drier contexts would decompose; e.g., wood or soft tissue as well as macro and microscopic ecofactual (i.e., organic without human workmanship, yet culturally relevant) material⁶¹, and now even ancient molecules including DNA^{62,63}. Thus, efficient drainage via ditches represents a direct threat to the survival of that material, including in some cases, the ditch itself. Similarly, palaeoecologists have usually understood drainage ditches to have generally negative impacts on the preservation of palaeoecological records although, if abandoned and




Fig. 2 | Drainage ditches can give clues to lost landscapes. Ditches in the Netherlands (52.20°N 5.12°E) show a wedge-shaped structure, and point towards the top of the (now missing) dome of a destroyed raised bog. Maps data: Google Earth, Maxar Technologies.

silted-up, they can provide excellent ecological records spanning centuries to millennia^{64–66}.

Ditches have left an archaeological and historic legacy of past humans and their relationships to the wider environment, as well as a paper archive where ditches have been mapped and described. These records attest to the fact that many regions' wetland environments can be more a product of ditches (irrigation and drainage), and thus human agency, rather than of geology and climate. In drained contexts, the outcome has been described as a 'reclamation landscape'⁶⁷. Key challenges moving forward will be to understand how these histories of reclamation or irrigation have impacted an array of different ecosystem services, and how this combined origin then frames and directly impacts drivers for wetland restoration. There is a possibility that without a full understanding of the deeper history of ditches in any given location restoration activities (blocking ditches or reducing irrigation) might have perverse outcomes. For example, the draining of swamp and marshland has generally been associated with the historical reduction in mosquito-borne diseases (such as malaria and Dengue Fever). Here, a clear understanding of vector ecology, as well as educational efforts, may be required to balance the pros and cons of wetland restoration in the public imagination⁶⁸. Given the overlapping, multi-directional, and complex network of ecosystem services that are underpinned by ditches, holistic inter- and transdisciplinary approaches will be required when making decisions relating to the tradeoffs involved regarding changes to ditch hydrological regimes. Recognition of the importance of ditches as socio-ecosystems is an important first step.

Generally, active ditches contribute to human wellbeing through ecosystem services across all categories: cultural, regulating, and provisioning (supporting ecosystem services are often left out of current frameworks^{69,70}) (see Supplementary Table 1). Traditional ditch systems—generally gravity-fed, not lined or covered—often supply water to relatively small-scale agriculture, and as such are sometimes seen as low-value or somewhat primitive. Fernald et al.⁷¹ examined hydrological ecosystem services in irrigation systems in New Mexico, USA, traditionally referred to as

“acequias”. Depending on soil type and gradient, these traditional irrigation systems can yield later-season return flows to mainstem rivers. This service is crucial in arid environments with flashy hydrographs. Raheem et al.⁷² examined a comprehensive suite of ditch-related ecosystem services for traditional irrigating communities in northern New Mexico, USA. A significant contribution of that work is to use the traditional Spanish terminology from those communities to describe a range of landforms usually described in English in the ecological literature. This sort of translation, from conventional scientific or planning terminology to local usage, is crucial in ascertaining the extent of cultural ecosystem services⁷³. Others have looked at ditches through an ecosystem services lens; for example, when considering sediment retention and bird habitat in northern Mexico⁷⁴, investigating the ‘multifunctionality’ of agricultural water use⁷⁵, and the “nature-based solutions” provided by subak irrigation systems in Bali⁷⁶.

Many ditches, and particularly irrigation channels, are designed to provide provisioning ecosystem services. Ditches carry water to farm fields, to orchards, to arbors, and to fish ponds to support food production systems for human use. Ditches are used to produce power, by driving mills of various sorts⁷⁷, which in turn help to produce food and goods. Sometimes fish, crustaceans^{78,79}, and other species that inhabit ditches provide nourishment for humans. Humans and animals drink water from ditches, although many ditches carry away waste and sewage⁸⁰. This latter function, both historically and to the present day, can have negative impacts on people who live within ditched landscapes or urban spaces. Indeed, ditches can and do often deliver an array of ecosystem disservices many of which (e.g., pollutant dispersal, greenhouse gas emission, facilitation of the movement of invasive species) are discussed later. Furthermore, even when ditch construction creates certain beneficial ecosystem services, other ecosystem services can be lost due to the associated land drainage (e.g., loss of wetlands and their carbon sink capacity¹⁰) or degradation of natural stream channels (e.g., straightening of channels and removal of riparian zones which would otherwise reduce peak flows and lead to less flashy systems¹¹).

There are also many examples of cultural ecosystem services arising from ditches, including traditional practices around the governance of irrigation systems around the world^{77,81–83}. Festivals and other occasions may be timed around irrigation calendars, and annual ditch cleanings, blessings of irrigation waters, and harvest festivals often play a large role in many communities⁷². These services can also include traditions that arise in irrigation systems and communities that are not directly related to governance, such as the Matachines dances in northern New Mexico, USA, or stories about mythical beings (e.g., La Llorona in Mexico, a ghost that resides near waterbodies). These cultural ecosystem services often have a deep history and are related to many of the systems previously described with regard to archaeology.

Ditches can bring people together in some of the best and some of the worst ways. Beyond celebrations, the need for neighbourly cooperation in ditch maintenance has been common throughout history and was addressed in Swedish medieval provincial laws (*landskapslag*)⁸⁴, practiced among villagers in Tokugawa Japan⁸⁵ and continues today in many countries^{86,87}. In the USA, such cooperation may take the form of authorities known as drainage districts or water management districts, whilst the UK has Internal Drainage Boards. Contrastingly, in contemporary legislation, ditches may be mostly ignored, e.g., in the USA Clean Water Act⁸⁸, and in New Zealand regulations where livestock do not have to be fenced out of agricultural ditches and streams <1 m wide and <30 cm deep⁸⁹. When this lack of regulation is combined with their capacity to carry harmful pollutants and pathogens, whose load is typically elevated, often deliberately concentrated, in areas of high poverty⁹⁰, then artificial channels become hotspots for environmental injustice and exploitation^{13,91,92}. Historically, drainage and irrigation have even served as tools of colonization, impinging upon the swampy refuges of indigenous and enslaved communities⁹³, and “reclaiming” arid environments for Western-style agriculture⁹⁴. Ditches can exist not just as public/municipal or private/individual infrastructure, but often at the nexus of the two, which can be pushed in either direction along the ditch continuum in exercises of power. In underserved communities

today, if there is stormwater infrastructure at all, privately maintained ditches may take the place of public, more expensive subsurface infrastructure. Ditches’ very existence can become a symbol of neglect and disempowerment^{95–97}. Yet, local understanding of what standard public maintenance would achieve socially and environmentally (e.g., improved downstream water quality^{98,99}) may fall short of the science. A better understanding, and regulatory acknowledgement, of ditches could empower others to help redesign ditches to best suit their priorities.

Physical and hydrological

Ditches are found in diverse catchments, spanning natural (forests and wetlands)^{100,101} to intensively managed (urban and agricultural) land uses^{18,102,103} (Fig. 1). The morphology of ditches depends on the surrounding landscape and required function: drainage, irrigation, hydropower, transportation or boundary delineation (see Supplementary Table 1). Ditches often have relatively straight, narrow and deep channels (Fig. 1A, F). Floodplains may be disconnected, or absent entirely, and there is reduced lateral hydrological connectivity with the riparian zone. As such, ditches are recipients of terrestrial fluxes, often from intensively managed environments^{104,105}. Because gully processes rapidly degrade ditches with steep bed slopes, ditches are primarily a feature of low elevation gradient environments. Depending on their position in the hydrological network, ditches generally drain catchments to lower the groundwater table (e.g., to enable food and fiber production, including forestry and peat extraction), redirect water (e.g., in polder systems or fenland), serve as distributive irrigation systems, or serve as temporary water storage locations which can help mitigate impacts of peak flows and downstream flooding. Regardless of function, ditches alter the original water balance of a catchment and the downstream hydrological regime¹¹.

Similar to natural streams, ditch hydrological connectivity with surrounding terrestrial environments, and thus biogeochemical exchange, depends primarily on their local geography and geomorphology, i.e., position within the landscape, length, width, presence and density of subsurface drainage, and channel substrate which may be permeable (sand, loam, peat, Fig. 1C, D, F, H) or impermeable (clay, concrete)^{106,107} (Fig. 1B, E, G). Regardless of channel substrate, many ditches are subject to gradual sediment and organic matter accumulation both in the form of alluvium from upstream reaches, fine sediments brought in by the tide in coastal areas, in-situ degradation of plants, and via bank erosion^{108,109}, necessitating occasional dredging to maintain original hydraulic function^{110,111}. By providing alternative flow paths, ditches can help moderate erosion elsewhere in the landscape⁷².

Depending on the hydraulic gradient between ditches and surrounding land, through riparian and hyporheic zones, ditches can either gain or lose water, with consequences for their water balances and flow^{102,112}. Depending on the flow, ditches can include dry channels (Fig. 1C) that flow intermittently for irrigation or drainage, still or slow-moving waters, to fast-moving and flashy environments^{108,113}. Their hydrology is determined by their topographical position^{11,111}, being generally more dynamic in ditches positioned along existing flow pathways or running downslope, and more static in ditches positioned in lowlands between flow pathways or following contours. Ditches that follow existing flow pathways may be difficult to distinguish from straightened or modified streams, whereas ditches that do not follow existing flow pathways are artificially constructed waterways. Thus, the hydrological connectivity of ditches to the stream network and downstream aquatic ecosystems depends largely on their topographic setting and design, but can be modified by their specific purpose, management, and age (as abandoned ditches may become disconnected from the stream network). Whether a ditch is connected to a stream network has important implications for flow direction, the transport and magnitude of lateral chemical fluxes, and pollution risks for downstream ecosystems¹⁰⁹.

In addition to connecting to and often replacing portions of natural hydrological networks, ditches themselves are often installed in geometric networks that expand the hydrological network structurally and functionally (Fig. 3). Frequently these networks are associated with subsurface pipes, sluices, weirs, and pumps^{114–116}. These structures are used to control the



Fig. 3 | Ditch networks can be extensive, vary in form/arrangement, and occur in different land covers. A Oil palm plantation on drained peatlands in Sarawak, Indonesia (2.66°N, 112.45°E), B agricultural land in The Netherlands (51.89°N 4.83°E), C drainage and irrigation canals in the Mesopotamian Marshes, Iraq

(30.96°N, 46.94°E), D urban canals in Xochimilco, Mexico City (19.27°N 99.09°W). Maps data: Google Earth, Airbus, Maxar Technologies (A), Google Earth, Airbus (B, D), Google Earth, Airbus, Maxar Technologies, CNES / Airbus.

direction and volume of water flow, and can shunt flows between ditches and their inlets and outlets, thus bypassing buffering effects of soils, riparian habitats, and other natural ecosystem components. Ditches often receive water and pollutant fluxes from artificial drainage, such as urban storm-water or agricultural subsurface drains^{18,104,117}, or through direct water pumping into ditches¹¹². With human pumping, with or without pipe networks, ditches can also convey water uphill. Thus, ditches are much more likely than non-tidal natural linear channels to flow in both directions at least occasionally, even if designed primarily for either irrigation or drainage. Due to the ability of ditches to defy natural watershed boundaries, the land areas with which ditches interact hydrologically can prove challenging to delineate, which can hinder the upscaling of aquatic fluxes of carbon and other solutes¹¹⁸.

Another important aspect determining the hydrological regime of ditches and their connectivity to terrestrial and other aquatic ecosystems, is their management²⁰. At the most basic level a distinction can be made between drainage ditches, which lower the adjacent terrestrial water table, and irrigation channels, which can contribute significantly to groundwater recharge via irrigation return flow¹¹⁹. Management can include channelization, dredging, vegetation removal, flow and water-level regulation^{111,118,120}, or riparian zone management (e.g., vegetation removal, establishing trees, widening flood-plains, fencing out livestock^{104,121}), all of which have the potential to affect in-situ hydrological processes as well as the eco-hydrology of the surrounding landscape. For example, the removal of vegetation and accumulated sediment from boreal forest ditches is periodically done to lower water tables to improve productivity, but it also results in increased concentrations of suspended sediment and nutrients¹²². Conversely, fencing out livestock from agricultural ditches has been shown to reduce bank erosion and thus decrease downstream nutrient and sediment loads¹²¹. Many types of management are deployed in combination, and integrated studies are needed to fully understand their cumulative impacts on ecohydrology.

As flow regimes in ditches change seasonally following either natural hydrological cycles or human activities¹²³, water retention time in ditches can

vary dramatically, with consequences for biogeochemical function^{11,109,124}. Due to their small size and volume, ditches are usually very sensitive to changes in flow, and can periodically be dry or overflowing. Thus, with more extreme hydrological events and changing global climate, many existing ditches may require adjustments in size, design, management, spatial density and/or expectations of function. The ultimate physical challenge of climate change will require careful consideration and balancing of multiple ecosystem services moving forward.

Biogeochemical

Ditches play a unique biogeochemical role due to their high connectivity with the surrounding landscape, processing inputs of nutrients, sediments, and pollutants from the wider environment¹⁰³. Ditches, like other aquatic systems, are not passive pipes but active components of global biogeochemical cycles¹²⁵, and the biogeochemical fingerprint of water leaving a ditch system can be substantially altered relative to incoming flow. The degree of biogeochemical processing is controlled mainly by residence time (the amount of time water spends in the ditch before flowing elsewhere), as observed in other inland waters^{126,127}. Residence time is itself determined by size, hydrological flow conditions, vegetation, landscape setting (e.g., upland versus lowland, effective catchment area), and management regime. Despite the global prevalence of ditches, they remain neglected in contemporary syntheses of the role of aquatic environments in biogeochemical cycles^{128–130}. Omitting ditch processes from large-scale biogeochemical assessments, including global carbon models, could lead to significant errors. Here, we propose that ditches are best viewed as ubiquitous reactive surfaces; as lines that bind across and beyond catchments.

Ditches can act as nutrient sinks or sources within the landscape, and this dual functionality is dependent on multiple factors including nutrient load, flow direction, vegetation cover and type, and climate. Therefore, sink or source behaviour can vary significantly over space and time¹¹⁷. As in other aquatic ecosystems, high nutrient loading can cause a wide range of problems, including excess phytoplankton growth, declines of aquatic

macrophytes, loss of biodiversity, and oxygen depletion¹³¹. These eutrophication effects are especially common in drainage ditches¹³² because they often serve as headwaters that receive anthropogenic terrestrial nutrient loads first before transferring them to receiving waters, including, rivers, wetlands, lakes, seas¹³³, and groundwater¹³⁴. Similarly, irrigation ditches can convey nutrients up gradient to land and municipal water systems¹³⁵.

Similar to other freshwater environments, vegetation and water residence time are important drivers of nutrient removal capacity^{124,136,137}. Longer residence times (which may be facilitated by water control structures), contribute to the creation of anaerobic conditions, promote organic matter accumulation, and encourage microbial denitrification^{104,138}. Reactive nitrogen (N) can also be removed by plant uptake^{137,139,140}. Thus, ditches can mitigate N-loading from both agricultural activity¹³⁶ and urban runoff, often at rates comparable to or exceeding those of natural systems^{141,142}. Vegetative uptake and sediment storage are also important pathways for phosphorus (P) removal in ditches and these processes, given time, can sometimes reverse even intense eutrophication¹⁴³. Recovery timescales from nutrient and other forms of pollution are often difficult to predict because of legacy effects, whereby nutrients stored in sediments can be mobilized under certain physical and chemical conditions^{144–146} or stored long-term, resulting in slow response times to changed nutrient loads^{132,147}. For example, practices such as dredging can destabilize banks and reduce P sorption capacity, resulting in ditches becoming a P source rather than sink¹⁴⁸.

Besides nutrients, ditches also receive runoff and groundwater comprising a complex mixture of chemical compounds. These mixtures may include microplastics¹⁴⁹, pathogens¹⁵⁰, antimicrobial resistant genes¹⁵¹, animal-borne hormones¹⁵², pharmaceuticals¹⁵³, trace metals¹⁵⁴, pesticides¹⁵⁵, and salts¹⁵⁶, which may, alone or in combination, have unintended and as yet largely unknown effects on ditch ecosystems. Roadside (Fig. 1E) and urban ditches (Fig. 1A), in particular, may receive a toxic mix of runoff that includes heavy metals, PAHs (polycyclic aromatic hydrocarbons), tire materials, pesticides, exhaust emissions, nutrients, road salt, and Per- and PFAS (polyfluoroalkyl substances- so called “forever chemicals”)^{157–159}. Despite this, ditch networks can be managed to retain pollutants in runoff^{160–162}. Indeed, because ditches often disproportionately contribute to catchment nutrient, sediment and pathogen loads⁸⁹ they could be prime candidates for pollutant removal and mitigation. Given the high degree of hydrological connectivity of ditches, water quality monitoring in these environments may yield important insights into emergent pollutant and contaminant pressures before they propagate to downstream environments.

The biogeochemistry of ditches extends beyond water quality, into landscape and global carbon and greenhouse gas (GHG) balances. The creation of ditches, especially in organic soils (Fig. 1D, F, H), causes adjacent terrestrial soils to dry out, which leads to extensive soil subsidence, oxidation, and carbon dioxide emissions¹⁶³. This drainage also affects concentrations of organic carbon, in particulate (POC) and dissolved organic carbon (DOC) forms, often leading to increases in the overall aquatic carbon flux¹⁶⁴. Increasing DOC concentrations may also pose additional problems, for example by imposing additional costs at drinking water treatment works¹⁶⁵.

An important fate of DOC and sediment organic matter in ditches is emission to the atmosphere in the form of GHGs. High loading of organic matter and nutrients, accumulation of organic sediment, fluctuating water table levels, connection to groundwater, slow water flow, low oxygen concentrations, and abundant vegetation (Fig. 1A) can render ditches as important sites for the production and release of the GHGs methane (CH₄), nitrous oxide (N₂O), and carbon dioxide (CO₂). The importance of ditches in drained organic soils (Fig. 1D, F, H) as landscape-scale hotspots of CH₄ emission has been known for decades¹⁶⁶. This arises because drainage effectively stops CH₄ emissions from adjacent terrestrial soils, leaving ditches as hotspots of CH₄ emission in the landscape. Recent evidence has shown this hotspot effect can extend to regional and national scales, and applies to mineral soils too^{22,167}. Ditches contribute significantly to global CH₄ emissions, contributing 3.5 Tg CH₄ yr⁻¹; equivalent to 1% of global anthropogenic CH₄ emissions⁹. In recognition of the potential for high area-

specific emissions, the Intergovernmental Panel on Climate Change (IPCC) now has guidelines for reporting of ditch CH₄ emissions¹⁶⁸.

Ditches can act as sources and sinks for CO₂. Carbon uptake by within-ditch vegetation, or the settling of suspended particulates, can lead to carbon accumulation in sediments^{169,170}, although this sediment is vulnerable to management interventions and storms¹⁷¹. There is a lack of knowledge about whether ditches also act as landscape-scale hotspots of CO₂ emission. Some evidence suggests that the CO₂ emissions from ditches may be small relative to those of the terrestrial ecosystem they drain^{164,172} (but see Hendriks et al.¹⁷³). Indeed, blocking ditches to restore water tables and thereby reduce CO₂ emissions is a key focus for land management in areas dominated by organic soils, which can lead to either the removal of ditches¹⁷⁴, or their modification to act as (effectively) irrigation rather than drainage systems. By altering overall water residence times, drainage networks potentially enhance rates of organic matter processing within freshwater ecosystems, and thereby reduce the organic carbon flux from land to ocean¹⁷⁵. The resulting CO₂ emissions from ditch surfaces are rarely measured, and could represent a missing term in landscape or catchment-scale carbon budgets¹⁷⁶. At the global scale, ditch CO₂ emissions are estimated at 30 Tg C yr⁻¹¹⁷⁷.

Nitrogen-enriched water discharging into ditches is well recognized as a source of indirect N₂O emissions in agricultural landscapes^{178–180}, and IPCC methodologies have been developed to facilitate reporting of these emissions¹⁸¹. Still, fewer studies have measured N₂O emissions from ditches directly. In a recent review, Webb et al.¹⁸² detected significant spatial and temporal variation in emissions between different types of artificial waters including ditches. This variability is only partially explained by nitrate loading from surrounding catchments (e.g., N application to agricultural fields), as assumed in IPCC reporting, indicating additional drivers¹⁸³. Although there is continued uncertainty in the magnitude and timing of ditch N₂O emissions, and their importance at landscape scales (i.e., when compared to terrestrial emissions), the global ditch emission is 0.03 Tg N yr⁻¹¹⁷⁷.

Floral and faunal

The ecology of ditches has been overlooked from both research and conservation perspectives. Many ditches are surrounded by agriculture (Fig. 1D, F, H) or urban landscapes (Fig. 1A, B, E, Supplementary Table 1) and heavily managed to facilitate a particular hydrological regime, roughness, or vegetation cover. Despite this, many ditches represent novel ecosystems with distinct biotic communities^{21,184–186}, and can support a high biodiversity of aquatic macroinvertebrates^{106,187}, waterbirds^{188,189}, amphibians¹⁹⁰, macrophytes^{191,192} and fish^{193,194}. Ditches can also support floral and faunal assemblages similar to adjacent natural streams^{123,195}, lakes⁷, and wetlands¹⁷⁰, and can be used by fish, amphibians, turtles, plants, and other taxa for reproduction^{12,196,197}. At a landscape scale, the unique hydrological characteristics of ditches can provide important ecological niches for many species, thereby increasing regional biodiversity^{186,195,198,199}. However, habitat quality and the biodiversity value of ditches is heavily dependent on management regimes^{134,200}. Ditches with a variety of successional stages (environmental heterogeneity) and good water quality, that enables diverse aquatic plant communities to develop, tend to support the most diverse fauna^{12,186,198}. Conversely, in ditches where riparian vegetation is periodically removed, steep bank angles maintained via dredging, or nutrient levels elevated, ditch freshwater biodiversity can be amongst the lowest in the wider landscape^{201,202}. To maintain a ditch's original function regular management is necessary. However, where this can be undertaken with a consideration for biodiversity (e.g., marginal vegetation managed on one bank, macrophytes maintained in selected areas or habitat diversity increased^{106,187,201}), ditches can display a very high conservation value²⁰³.

In intensively farmed or urbanized landscapes (Fig. 1A, B, E, F, H, Supplementary Table 1), ditches often represent the only available refugia for aquatic and riparian wildlife and can reflect a range of natural habitat analogues that have been lost due to agricultural intensification, land drainage, and water abstraction^{123,204}. As ditches can possess slow flow, still water, and highly variable water depths, ditch networks often support species that are typical of both lotic and lentic habitats^{187,205}. However, the extent and

importance of this “refugia” function depends on management regimes and the type of ditch system (e.g., concrete-lined systems will be less desirable for fish, boat traffic can lower plant biodiversity¹²), as well as wider catchment characteristics such as the presence of other waterbodies or set aside land²¹. Where ditches are located within and close to ancient wetland habitats, they may act as a ‘memory’ or palimpsest of lost aquatic habitats, and continue to support remnant flora and fauna present in lost natural lentic, river, fen and marshland habitats²⁰⁶. Evidence from the Fens of East Anglia (UK) show a high degree of correspondence between the freshwater flora of fenland farm ditches and that of Wicken Fen (the UK’s oldest nature reserve)²⁰⁷ and a lesser, but nevertheless strong, representation of the historic fen and marsh species. Ditches often support rare species (e.g., gastropods²⁰⁸) and one of the UK’s rarest plant species, the fen ragwort (*Senecio paludosus*), survives in one Fenland ditch²⁰⁹.

Ditches also provide connectivity between natural aquatic habitats through an often hostile terrestrial matrix (*sensu* Mazerolle²¹⁰). Plants, amphibians, fish, crayfish, and turtles have been shown to use ditches to disperse across the landscape, highlighting how ditches can act as connectivity corridors and sometimes mitigate negative effects of habitat fragmentation^{12,197,211}. Ditches in intensively managed landscapes may also provide habitat and passage for fully terrestrial species²¹². Inevitably this connectivity can also aid the dispersal of invasive and exotic species^{114,213,214}, which is an issue most ditch managers have yet to consider. Ditches not only provide hydrological connectivity but contribute important aquatic-terrestrial linkages. For example, ditch invertebrate communities transfer energy to the riparian zone and provide trophic resources for mammals and birds²¹⁵, and critical regulating ecosystem services including habitat for a reservoir of pollinators²¹⁶. In this way ditches, comprising an aquatic habitat, bankside vegetation, and potentially with an uncultivated headland on one or both sides, offer the potential to establish ‘blue-green corridors’ of biodiversity reflecting both wetland and terrestrial biota and providing reservoirs of natural predators of crop pests that could provide multiple environmental services in arable landscapes. Such ‘linear nature hotspots’ with their close juxtaposition of different ecotones would require less land to be taken out of production than area-based conversion of fields for a similar gain in biodiversity.

Synthesis and conclusions

We have demonstrated that ditches are important socio-ecosystems³⁰ whereby human society and nature are interconnected and co-evolving. Ditches exert a range of effects upon adjacent environments across scales, and have done so for at least the past 8000 years. These effects can be both positive (e.g., support for global crop production and thus the whole of human society, refuges for biodiversity, processing pollutants, sites of important cultural ecosystem services) and negative (e.g., GHG emissions, dispersal of pollutants downstream, degradation of biodiversity, damaging palaeoecological records, habitats for invasive species). An emerging challenge is to manage these waterways in a multifunctional manner; that is, maximizing positive synergies whilst minimizing trade-offs, in order to deliver multiple ecosystem services at the same time. This is difficult, with no “one size fits all” solution, because of the many different social and ecological purposes that ditches are managed for (Fig. 1, Supplementary Table 1). Some opportunities for multifunctionality have been demonstrated; e.g., raising ditch water levels in temperate agricultural peatlands during the winter season has been shown to decrease carbon dioxide emissions without any negative effects on crop yield²¹⁷; carefully designed riparian integrated buffer zones can reduce ditch nutrient loads, increase biodiversity, and allow the production of biomass²¹⁸. Nevertheless, in many cases the primary function of a human-created channel may be incompatible with secondary co-benefits¹³, e.g., irrigation and urban channels built to transport water may be dredged of sediment and vegetation to increase water flows, which therefore vastly reduces biodiversity and aesthetic value. Further difficulties in managing for multifunctionality arise because ditches frequently cross land boundaries and catchments, with different owners/managers being responsible for consecutive sections of channel. Collaborative thinking is

required between individuals or organizations if a management strategy is to be coherent, effective²¹⁹, and environmentally just. Finally, global and climatic change will increasingly stress the functioning of ditches, and in many ways human reliance on them for water security will increase as climate patterns and flow regimes become more unpredictable. Therefore, ditches will require ongoing adaptive management²⁶. Considering these disparate and diverse issues, ditch management can clearly be perceived as a “wicked problem”²²⁰.

Despite this complexity, there are opportunities in ditch management, because ditches are extensive, important, undervalued, and understudied. Additionally, because ditched landscapes are often intensively managed makes them excellent candidates for testing novel environmental management strategies¹⁸. Many channels are overbuilt for their intended purpose²²¹ (e.g., roadside ditches may only rarely carry water during storms) and therefore slight changes to design or management could provide bonus ecosystem services. Even lack of regulation can present an opportunity to innovate. However, questions arise when considering ditches through the framework of ecological concepts such as conservation and restoration: how do we apply these concepts to highly managed and artificial ditch ecosystems? The EU’s Water Framework Directive (WFD) can perhaps give some answers. Although ditches are generally overlooked in the WFD¹⁶, the WFD does provide guidance for heavily modified and artificial waterbodies. Specifically, the WFD acknowledges that these waterbodies may have important functions (e.g., navigation, land drainage, water regulation, etc) which may prevent them from achieving “good ecological status”. Instead, the aim is for these modified waterbodies to reach the lower threshold of “good ecological potential”²²².

Perhaps a creative rethinking of traditional paradigms in environmental management is necessary to answer such questions of conservation and restoration. Because of the aforementioned conflicts in management at the ditch-scale, it may be that the opportunity for multifunctional management only arises when ditches, in conjunction with their associated connections to groundwater, wetlands, streams and lakes, are considered as habitat networks²²³ or meta-ecosystems (*sensu* Loreau et al.²²⁴) at the landscape scale; when entire ditch networks are adaptively managed for multiple social and environmental gains (as suggested for constructed wetland management²²⁵). Indeed, the necessity of landscape-scale thinking is now reflected in terms such as hydroscares²²⁶, pondscares²²⁷, riverscares²²⁸, wetlandscapes²²⁵, and wetscapes²²⁹, and “ditchscapes” should also be integrated into thinking about, and managing for, the wider environment. Another consideration is the temporal scales, which can dramatically alter how traditional biological conservation view ditches and the resulting ‘scapes’ they create²³⁰. The role of archaeology, history and allied disciplines is vital to understanding the deeper historical ditchscape trajectories for both hydrological and biodiversity management, but also cultural heritage, tangible and intangible.

The success of managing novel ditchscapes for multifunctionality depends on a variety of factors, but we believe that one of the most significant limitations is mapping. Simply put, we do not know where ditches are. Detailed channel maps, across a range of spatial scales, would help to solve an array of management questions (e.g., where do we prioritize ditch management?) and research gaps (e.g., how do we accurately upscale biogeochemical processes?). Mapping methods are constantly being refined and it is now possible to map ditches at local and regional scales using remote sensing and novel machine learning methods^{231–235}. However, location alone is only part of the puzzle because most purposes would also require information on ditch size (width and depth) and flow (perennial, intermittent, or ephemeral; see Fritz et al.²³⁶), at minimum. Channel width is essential to calculate ditch surface area, and is key for some global upscalings of processes and cycles as well as for IPCC reporting¹⁶⁸, yet it is frequently unknown⁹. We are optimistic that these issues will be resolved in the near future by the use of remote sensing based mapping aided by artificial intelligence methods⁸, provided that mapping efforts use these tools to include, rather than exclude, ditches¹⁷⁰.

A final challenge is the need to widely reframe how individuals, and society at large, perceive ditches. Assuming that artificial aquatic ecosystems

lack ecological value promotes neglect¹³. Environmental decision-makers assume ditches are low-quality ecosystems, and manage accordingly; subsequently, their assumption becomes a self-fulfilling prophecy. Considering the prevalence of English-language ditch idioms with negative connotations, it is clear that changing perceptions will be no easy task.

If the challenges are addressed, and opportunities seized, then multifunctional management of ditchscapes could be an effective way to boost ecosystem services and provide nature-based solutions at scale, and across a gradient of land cover intensities, from urban, through agricultural, to forest, and wetlands. Such successful ditchscapes would link to multiple Sustainable Development Goals including “clean water and sanitation”, “climate action”, and “life on land”²⁷. Ditches have the potential to serve as resilient waterways on a changing planet; lines in the landscape recording the past, providing for the present, and directing us onwards to a more sustainable future.

Received: 28 February 2025; Accepted: 13 August 2025;

Published online: 22 August 2025

References

1. Tamburrino, A. Water technology in Ancient Mesopotamia in *Ancient Water Technologies* (ed Mays, L.) 29–51 (Springer Netherlands, 2010).
2. Hassan, F. A. The dynamics of a riverine civilization: a geoarchaeological perspective on the Nile Valley, Egypt. *World Archaeol.* **29**, 51–74 (1997).
3. Schultz, B., Thatte, C. D. & Labhsetwar, V. K. Irrigation and drainage. Main contributors to global food production. *Irrig. Drain. J. Int. Comm. Irrig. Drain.* **54**, 263–278 (2005).
4. Shoop, S. A., Hills, J. & Uberuaga, J. Maintenance and drainage guidance for the Scott Base Transition, Antarctica. ERDC/CRREL TR-14-25. (2014). <https://erdc-library.erdcdren.mil/jspui/handle/11681/5505>.
5. Castellano, M. J., Archontoulis, S. V., Helmers, M. J., Poffenbarger, H. J. & Six, J. Sustainable intensification of agricultural drainage. *Nat. Sustain.* **2**, 914–921 (2019).
6. Paavilainen, E. & Päivänen, J. *Peatland Forestry: Ecology and Principles* Vol. 111 (Springer Science & Business Media, 1995).
7. Verdonchot, R. C., Keizer-vlek, H. E. & Verdonchot, P. F. Biodiversity value of agricultural drainage ditches: a comparative analysis of the aquatic invertebrate fauna of ditches and small lakes. *Aquat. Conserv. Mar. Freshw. Ecosyst.* **21**, 715–727 (2011).
8. Laudon, H. et al. Emerging technology can guide ecosystem restoration for future water security. *Hydrol. Process.* **36**, e14729 (2022).
9. Peacock, M. et al. Global importance of methane emissions from drainage ditches and canals. *Environ. Res. Lett.* **16**, 044010 (2021a).
10. Fluet-Chouinard, E. et al. Extensive global wetland loss over the past three centuries. *Nature* **614**, 281–286 (2023).
11. Bierzoza, M., Hallberg, L., Livsey, J., Prischl, L. A. & Wynants, M. Recognizing agricultural headwaters as critical ecosystems. *Environ. Sci. Technol.* **58**, 4852–4858 (2024).
12. Chester, E. T. & Robson, B. J. Anthropogenic refuges for freshwater biodiversity: their ecological characteristics and management. *Biol. Conserv.* **166**, 64–75 (2013).
13. Clifford, C. C. & Heffernan, J. B. Artificial aquatic ecosystems. *Water* **10**, 1096 (2018).
14. Saulnier-Talbot, É. & Lavoie, I. Uncharted waters: the rise of human-made aquatic environments in the age of the “Anthropocene”. *Anthropocene* **23**, 29–42 (2018).
15. Mace, G. M. Whose conservation? *Science* **345**, 1558–1560 (2014).
16. Biggs, J., Von Fumetti, S. & Kelly-Quinn, M. The importance of small waterbodies for biodiversity and ecosystem services: implications for policy makers. *Hydrobiologia* **793**, 3–39 (2017).
17. Hobbs, R. J., Higgs, E. & Harris, J. A. Novel ecosystems: implications for conservation and restoration. *Trends Ecol. Evol.* **24**, 599–605 (2009).
18. Tank, J. L., Speir, S. L., Sethna, L. R. & Royer, T. V. The case for studying highly modified agricultural streams: farming for biogeochemical insights. *Limnol. Oceanogr. Bull.* **30**, 41–47 (2021).
19. Koetsier, P. & McCauley, L. M. An irrigation canal as a lotic mesocosm: examining the relationship between macroinvertebrate benthos and drift. *West. North Am. Nat.* **75**, 259–270 (2015).
20. Dollinger, J., Dagès, C., Bailly, J. S., Lagacherie, P. & Voltz, M. Managing ditches for agroecological engineering of landscape. A review. *Agron. Sustain. Dev.* **35**, 999–1020 (2015).
21. Herzon, I. & Helenius, J. Agricultural drainage ditches, their biological importance and functioning. *Biol. Conserv.* **141**, 1171–1183 (2008).
22. Koschorreck, M. et al. Hidden treasures: human-made aquatic ecosystems harbour unexplored opportunities. *Ambio* **49**, 531–540 (2020).
23. Wu, W. et al. Agricultural ditches are hotspots of greenhouse gas emissions controlled by nutrient input. *Water Res.* **242**, 120271 (2023).
24. McMahan, E. A., Cloud, J. M., Josh, P. & Scott, M. Nature with a human touch: Human-induced alteration negatively impacts perceived naturalness and preferences for natural environments. *Ecopsychology* **8**, 54–63 (2016).
25. Everhart, T. D. On the monumentality of ditches. *J. Anthropol. Archaeol.* **62**, 101295 (2021).
26. Ross, M. R., Bernhardt, E. S., Doyle, M. W. & Heffernan, J. B. Designer ecosystems: incorporating design approaches into applied ecology. *Annu. Rev. Environ. Resour.* **40**, 419–443 (2015).
27. United Nations. 2015. Transforming our world: the 2030 Agenda for Sustainable Development. United Nations: New York, NY, USA.
28. Garcia, P. & Santistevan, M. Acequia waters: community resource or commodity? in *Water for the People: The Acequia Heritage of New Mexico in a Global Context* (eds Lamadrid, E. R., & Rivera, J. A.) (University of New Mexico Press, 2023).
29. Rivera, J. A. *Acequia Culture: Water, Land, and Community in the Southwest* (University of New Mexico Press, 1998).
30. Glaser, M., Ratter, B. M., Krause, G. & Welp, M. New approaches to the analysis of human–nature relations. in *Human-Nature Interactions in the Anthropocene* (3–12) (Routledge, 2012).
31. Wittfogel, K. A. *Oriental Despotism: A Comparative Study of Total Power* (Yale University Press, 1957).
32. Butzer, K. Irrigation, raised fields and state management: Wittfogel redux. *Antiquity* **70**, 200–204 (1996).
33. Oates, D. & Oates, J. Early irrigation agriculture in Mesopotamia. in *Problems in Economic & Social Archaeology* (ed Sieveking, G) 109–135 (Duckworth, 1976).
34. Hritz, C. & Wilkinson, T. J. Using Shuttle Radar Topography to map ancient water channels in Mesopotamia. *Antiquity* **80**, 415–424 (2006).
35. Rost, S. Water management in Mesopotamia from the sixth till the first millennium BC. *Wiley Interdiscip. Rev. Water* **4**, e1230 (2017).
36. Schrakamp, I. Irrigation in 3rd millennium southern Mesopotamia: cuneiform evidence from the Early Dynastic IIIb City-State of Lagash (2475–2315 BC). in *Water Management in Ancient Civilizations* (ed Berking, J.) (Topoi, 2018). <https://doi.org/10.17171/3-5>.
37. Storozum, M. et al. Early evidence of irrigation technology in the North China Plain: geoarchaeological investigations at the Anshang site, Neihuang County, Henan Province, China. *Geoarchaeology* **33**, 143–161 (2018).
38. Brohier, R. L. *Ancient Irrigation Works in Ceylon* (Government Publications Bureau, 1934).
39. Palacios-Fest, M. R. Nonmarine ostracode shell chemistry from ancient Hohokam irrigation canals in central Arizona: a paleohydrochemical tool for the interpretation of prehistoric human

- occupation in the North American Southwest. *Geoarchaeology* **9**, 1–29 (1994).
40. Simms, S. R., Rittenour, T. M., Kuehn, C. & Cannon, M. B. Prehistoric irrigation in Central Utah: chronology, agricultural economics, and implications. *Am. Antiquity* **85**, 452–469 (2020).
41. Lucero, L. J. The collapse of the Classic Maya: A case for the role of water control. *Am. Anthropol.* **104**, 814–826 (2002).
42. Goodbred, S. L. Jr, Dillehay, T. D., Mora, C. G. & Sawakuchi, A. O. Transformation of maritime desert to an agricultural center: Holocene environmental change and landscape engineering in Chicama River valley, northern Peru coast. *Quat. Sci. Rev.* **227**, 106046 (2020).
43. De Meulemeester, J. Granaries and irrigation: archaeological and ethnological investigations in the Iberian peninsula and Morocco. *Antiquity* **79**, 609–615 (2005).
44. Crifasi, R. *A Land Made from Water: Appropriation and the Evolution of Colorado's Landscape, Ditches, and Water Institutions* (University of Colorado Press, 2015).
45. Thompson, V. D. et al. Ancient engineering of fish capture and storage in southwest Florida. *Proc. Natl. Acad. Sci. USA* **117**, 8374–8381 (2020).
46. Gillmore, G. K. et al. Irrigation on the Tehran Plain, Iran: Tepe Pardis – The site of a possible Neolithic irrigation feature?. *Catena* **78**, 285–300 (2009).
47. Wilkinson, T. J., Rayne, L. & Jotheri, J. Hydraulic landscapes in Mesopotamia: The role of human niche construction. *Water Hist.* **7**, 397–418 (2015).
48. Sivapalan, M. & Blöschl, G. Time scale interactions and the coevolution of humans and water. *Water Resour. Res.* **51**, 6988–7022 (2015).
49. Maw, P., Wyke, T. & Kidd, A. Canals, rivers, and the industrial city: Manchester's industrial waterfront, 1790–1850. *Econ Hist. Rev.* **65**, 1495–1523 (2012).
50. Downward, S. & Skinner, K. Working rivers: the geomorphological legacy of English freshwater mills. *Area* **37**, 138–147 (2005).
51. Davies, H., Fyfe, R. & Charman, D. Does peatland drainage damage the palaeoecological record?. *Rev. Palaeobot. Palynol.* **221**, 92–105 (2015).
52. McCorvie, M. R. & Lant, C. L. Drainage district formation and the loss of Midwestern wetlands, 1850–1930. *Agric. Hist.* **67**, 13–39 (1993).
53. Chisholm, M. Navigation and the seventeenth-century draining of the Fens. *J. Hist. Geogr.* **32**, 731–751 (2006).
54. Ash, E. *The Draining of the Fens: Projectors, Popular Politics, and State Building in Early Modern England* (Johns Hopkins University Press, 2017).
55. Williamson, T. The disappearance of malaria from the East Anglian Fens. *Int. J. Region. Local Stud.* **2**, 109–122 (2006).
56. De Bont, C. H. M. *Vergeten Land: Ontginning, Bewoning en Waterbeheer in de Westnederlandse Veengebieden 800–1350* (Wageningen University, 2008).
57. Lewin, J. Enlightenment and the GM floodplain. *Earth Surf. Process. Landf.* **38**, 17–29 (2013).
58. Brown, A. G. et al. Natural vs anthropogenic streams in Europe: History, ecology and implications for restoration, river-rewilding and riverine ecosystem services. *Earth Sci. Rev.* **180**, 185–205 (2018).
59. Matthiesen, H., Brunning, R., Carmichael, B. & Hollesen, J. Wetland archaeology and the impact of climate change. *Antiquity* **96**, 1412–1426 (2022).
60. Van de Noort, R., Fletcher, W., Thomas, G., Carstairs, I. & Patrick, D. Monuments at risk in England's wetlands. Report for English Heritage (2002).
61. Gearey, B. & Chapman, H. *An Introduction to Peatland Archaeology and Palaeoenvironments* (Oxbow, 2022).
62. Cappellini, E. et al. Ancient biomolecules and evolutionary inference. *Annu. Rev. Biochem.* **87**, 1029–1060 (2018).
63. Brown, A. G. et al. New integrated molecular approaches for understanding lake settlements in NW Europe. *Antiquity* **96**, 1179–1199 (2022).
64. Horrocks, M. & Barber, I. Microfossils of introduced starch cultigens from an early wetland ditch in New Zealand. *Archaeol. Ocean.* **40**, 106–114 (2005).
65. Brown, A. G. et al. Late Quaternary evolution of a lowland anastomosing river system: geological-topographic inheritance, non-uniformity and implications for biodiversity and management. *Quat. Sci. Rev.* **260**, 106929 (2021).
66. Tol, G. et al. There's more than meets the eye": developing an integrated archaeological approach to reconstruct human–environment dynamics in the Pontine marshes (Lazio, Central Italy). *Geoarchaeology* **36**, 109–129 (2021).
67. de Haas, T. & Schepers, M. Wetland reclamation and the development of reclamation landscapes: a comparative framework. *J. Wetl. Archaeol.* **22**, 75–96 (2022).
68. Hawkes, F. M., Medlock, J. M., Vaux, A. G. C., Cheke, R. A. & Gibson, G. *Wetland Mosquito Survey Handbook: Assessing Suitability of British Wetlands for Mosquitoes* (Natural Resources Institute, 2020).
69. DeGroot, R. S., Wilson, M. A. & Boumans, R. M. J. A typology for the classification, description and valuation of ecosystem functions, goods and services. *Ecol. Econ.* **41**, 393–408 (2002).
70. Millennium Ecosystem Assessment (MEA). *Ecosystems and Human Well-Being: Synthesis* (Island Press, 2005).
71. Fernald, A. G., Baker, T. T. & Guldán, S. J. Hydrologic, riparian, and agroecosystem functions of traditional acequia irrigation systems. *J. Sustain. Agricult.* **30**, 147–171 (2007).
72. Raheem, N. et al. A framework for assessing ecosystem services in acequia irrigation communities of the Upper Rio Grande Watershed. *WIREs Water* **2**, 559–575 (2015).
73. Fleming, W. M., Rivera, J. A., Miller, A. & Piccarello, M. Ecosystem services of traditional irrigation systems in northern New Mexico, USA. *Int. J. Biodivers. Sci., Ecosyst. Serv. Manag.* **10**, 343–350 (2014).
74. Cital, F. et al. Ecosystem services (ES) provided by ditches in a desert agricultural valley. *Ecol. Eng.* **174**, 106462 (2022).
75. Groenfeldt, D. Multifunctionality of agricultural water: looking beyond food production and ecosystem services. *Irrig. Drain. J. Int. Comm. Irrig. Drain.* **55**, 73–83 (2006).
76. Risna, R. A., Rustini, H. A., Buchori, D. & Pribadi, D. O. Subak, a nature-based solutions evidence from Indonesia. in *IOP Conference Series: Earth and Environmental Science* (Vol. 959, p. 012030) (IOP Publishing, 2022).
77. Glick, T. F. & Kirchner, H. Hydraulic systems and technologies of Islamic Spain: history and archaeology. in *Working with Water in Medieval Europe* (267–329) (Brill, 2000).
78. Sato, M., Kumagai, M. & Azuma, N. Life history of *Tanakia lanceolata* in irrigation channel network of the Iwaki river basin in Northern Japan. *Trans. Jpn. Soc. Irrig., Drain. Rural Eng.* **80**, 233–243 (2012).
79. Chen, R. S., Wang, K. L. & Wu, C. Y. A preliminary study on the improvement of the ecological corridor in paddy fields. *Irrig. Drain.* **64**, 115–123 (2015).
80. Moeder, M. et al. Potential of vegetated ditches to manage organic pollutants derived from agricultural runoff and domestic sewage: A case study in Sinaloa (Mexico). *Sci. Total Environ.* **598**, 1106–1115 (2017).
81. Lansing, J. S. Balinese “water temples” and the management of irrigation. *Am. Anthropol.* **89**, 326–341 (1987).
82. Araral, E. Jr. What explains collective action in the commons? Theory and evidence from the Philippines. *World Dev.* **37**, 687–697 (2009).
83. Ilahiane, H. The break-up of the Ksar: changing settlement patterns and environmental management in Southern Morocco. *Afr. Today* **48**, 21–48 (2001).

84. Jakobsson, E. Ditching from a water system perspective. Draining the Swedish water landscape 1200–1900. *Water Hist.* **5**, 349–367 (2013).
85. Satō, T. Tokugawa Villages and Agriculture (trans. Mikiso Hane). in *Tokugawa Japan: The Social and Economic Antecedents of Modern Japan* (trans. Conrad Totman) (University of Tokyo Press, 1990).
86. The Environment Agency. *Living on the Edge; A Guide to Your Rights and Responsibilities of Riverside Ownership* (The Environment Agency, 2014).
87. Hannah, C. *When the Canals Run Dry: New Institutions and the Collective Governance of Irrigation Systems in Tajikistan* PhD thesis. (Duke University, 2018). <https://dukespace.lib.duke.edu/dspace/handle/10161/17456>.
88. Doyle, M. W. & Bernhardt, E. S. What is a stream?. *Environ. Sci. Technol.* **45**, 354–359 (2011).
89. McDowell, R. W., Cox, N. & Snelder, T. H. Assessing the yield and load of contaminants with stream order: Would policy requiring livestock to be fenced out of high-order streams decrease catchment contaminant loads?. *J. Environ. Qual.* **46**, 1038–1047 (2017).
90. Horvath, I. R. et al. Volunteer science data show degraded water quality disproportionately burdens areas of high poverty. *J. Hydrol.* **613**, 128475 (2022).
91. Purifoy, D. M. North Carolina [Un] incorporated: Place, race, and local environmental inequity. *Am. Behav. Sci.* **65**, 1072–1103 (2021).
92. Flowers, C. C. Hurricane Ida shows the one-two punch of poverty and climate change. *Nature* **597**, 449–449 (2021).
93. Emanuel, R. E. Water in the Lumbee world: a river and its people in a time of change. *Environ. Hist.* **24**, 25–51 (2019).
94. Morita, A. Infrastructuring amphibious space: the interplay of aquatic and terrestrial infrastructures in the Chao Phraya delta in Thailand. *Sci. Cult.* **25**, 117–140 (2016).
95. Hendricks, M. D. *The Infrastructures of Equity and Environmental Justice*. Texas A&M Univ. (2017).
96. Seamster, L. & Purifoy, D. What is environmental racism for? Place-based harm and relational development. *Environ. Sociol.* **7**, 110–121 (2021).
97. Watkins, K. in *Houston Public Media* (PBS NPR, 2022).
98. The Virginian Pilot. Pilot Online. Ditch drainage a lingering issue. in *Virginian Pilot* (2013). <https://www.pilotonline.com/2013/10/24/ditch-drainage-a-lingering-issue/>.
99. Bova, C. J. & Morrow, G. C. *Drowning a County: When Urban Myths Destroy Rural Drainage* (Random Tangent Press, 2014).
100. Hökkä, H., Stenberg, L. & Laurén, A. Modeling depth of drainage ditches in forested peatlands in Finland. *Baltic Forestry* 2020; 26, article ID 453. <https://doi.org/10.46490/BF453> (2020).
101. Stenberg, L. et al. Ditch erosion processes and sediment transport in a drained peatland forest. *Ecol. Eng.* **75**, 421–433 (2015).
102. Magner, J. et al. Channel evolution of Des Moines Lobe till drainage ditches in southern Minnesota (USA). *Environ. Earth Sci.* **67**, 2359–2369 (2012).
103. Needelman, B. A., Kleinman, P. J., Strock, J. S. & Allen, A. L. Drainage Ditches: improved management of agricultural drainage ditches for water quality protection: an overview. *J. Soil Water Conserv.* **62**, 171–178 (2007).
104. Hallberg, L., Hallin, S. & Bieroza, M. Catchment controls of denitrification and nitrous oxide production rates in headwater remediated agricultural streams. *Sci. Total Environ.* **838**, 156513 (2022).
105. Jia, Z. et al. Hydraulic conditions affect pollutant removal efficiency in distributed ditches and ponds in agricultural landscapes. *Sci. Total Environ.* **649**, 712–721 (2019).
106. Gething, K. J., Ripley, M. C., Mathers, K. L., Chadd, R. P. & Wood, P. J. The influence of substrate type on macroinvertebrate assemblages within agricultural drainage ditches. *Hydrobiologia* **847**, 4273–4284 (2020).
107. Liu, F. et al. Influence of substrates on nutrient removal performance of organic channel barriers in drainage ditches. *J. Hydrol.* **527**, 380–386 (2015).
108. Hallberg, L., Djodjic, F. & Bieroza, M. Does stream remediation reduce phosphorus and sediment exports from agricultural catchments?. *EGUosphere* **2023**, 1–24 (2023).
109. Moloney, T., Fenton, O. & Daly, K. Ranking connectivity risk for phosphorus loss along agricultural drainage ditches. *Sci. Total Environ.* **703**, 134556 (2020).
110. Aviles, R. D. B., Wesstrom, I. & Joel, A. A CSM-Based procedure for identifying segments of agricultural drainage ditches to be prioritized in maintenance work. *J. ASABE* **66**, 879–885 (2023).
111. Dollinger, J., Vinatier, F., Voltz, M., Dagès, C. & Bailly, J. S. Impact of maintenance operations on the seasonal evolution of ditch properties and functions. *Agric. water Manag.* **193**, 191–204 (2017).
112. Armstrong, A. & Rose, S. Ditch water levels manages for environmental aims: effects on field soil water regimes. *Hydrol. Earth Syst. Sci.* **3**, 385–394 (1999).
113. Strock, J. S., Kleinman, P. J., King, K. W. & Delgado, J. A. Drainage water management for water quality protection. *J. Soil Water Conserv.* **65**, 131A–136A (2010).
114. Griggs, P. D. Too much water: drainage schemes and landscape change in the sugar-producing areas of Queensland, 1920–90. *Aust. Geogr.* **49**, 81–105 (2018).
115. Page, S. et al. An assessment of the societal impacts of water level management on lowland peatlands in England and Wales: Report to Defra for Project SP1218: Managing agricultural systems on lowland peat for decreased greenhouse gas emissions whilst maintaining agricultural productivity. (UK CEH, 2020).
116. Valipour, M. et al. The evolution of agricultural drainage from the earliest times to the present. *Sustainability* **12**, 416 (2020).
117. Ezzati, G. et al. Impact of P inputs on source-sink P dynamics of sediment along an agricultural ditch network. *J. Environ. Manag.* **257**, 109988 (2020).
118. Holden, J. et al. The impact of ditch blocking on the hydrological functioning of blanket peatlands. *Hydrol. Process.* **31**, 525–539 (2017).
119. Jiménez-Martínez, J., Skaggs, T. H., Van Genuchten, M. T. & Candela, L. A root zone modelling approach to estimating groundwater recharge from irrigated areas. *J. Hydrol.* **367**, 138–149 (2009).
120. Aviles, D., Wesström, I. & Joel, A. Effect of vegetation removal on soil erosion and bank stability in agricultural drainage ditches. *Land* **9**, 441 (2020).
121. McKergow, L. A., Weaver, D. M., Prosser, I. P., Grayson, R. B. & Reed, A. E. Before and after riparian management: sediment and nutrient exports from a small agricultural catchment, Western Australia. *J. Hydrol.* **270**, 253–272 (2003).
122. Marttila, H. & Kløve, B. Dynamics of erosion and suspended sediment transport from drained peatland forestry. *J. Hydrol.* **388**, 414–425 (2010).
123. Carlson, E. A., Cooper, D. J., Merritt, D. M., Kondratieff, B. C. & Waskom, R. M. Irrigation canals are newly created streams of semi-arid agricultural regions. *Sci. Total Environ.* **646**, 770–781 (2019).
124. Webster, A. J., Groffman, P. M. & Cadenasso, M. L. Controls on denitrification potential in nitrate-rich waterways and riparian zones of an irrigated agricultural setting. *Ecol. Appl.* **28**, 1055–1067 (2018).
125. Cole, J. J. et al. Plumbing the global carbon cycle: integrating inland waters into the terrestrial carbon budget. *Ecosystems* **10**, 172–185 (2007).
126. Evans, C. D. et al. Variability in organic carbon reactivity across lake residence time and trophic gradients. *Nat. Geosci.* **10**, 832–835 (2017).
127. Jones, A. E., Hodges, B. R., McClelland, J. W., Hardison, A. K. & Moffett, K. B. Residence-time-based classification of surface water systems. *Water Resour. Res.* **53**, 5567–5584 (2017).

128. Drake, T. W., Raymond, P. A. & Spencer, R. G. Terrestrial carbon inputs to inland waters: A current synthesis of estimates and uncertainty. *Limnol. Oceanogr. Lett.* **3**, 132–142 (2018).
129. Gao, Y. et al. Carbon transportation, transformation, and sedimentation processes at the land-river-estuary continuum. *Fundam. Res.* **4**, 1594–1602 (2022).
130. Rosentreter, J. A. et al. Half of global methane emissions come from highly variable aquatic ecosystem sources. *Nat. Geosci.* **14**, 225–230 (2021).
131. Janse, J. H. & Van Puijenbroek, P. J. Effects of eutrophication in drainage ditches. *Environ. Pollut.* **102**, 547–552 (1998).
132. Van Puijenbroek, P. J. T. M., Cleij, P. & Visser, H. Aggregated indices for trends in eutrophication of different types of fresh water in the Netherlands. *Ecol. Indic.* **36**, 456–462 (2014).
133. Ahlgren, J., Djodic, F. & Löfgren, S. Åtgärder för att förbättra fosforretention i öppna diken i riskområden i jordbrukslandskapet runt Östersjön. En kunskapssammanställning. BalticSea2020, 33. (2011).
134. Webster, A. J. & Cadenasso, M. L. Cross-scale controls on the in-stream dynamics of nitrate and turbidity in semiarid agricultural waterway networks. *J. Environ. Manag.* **262**, 110307 (2020).
135. Mortensen, J. G. et al. Advancing the food-energy-water nexus: closing nutrient loops in arid river corridors. *Environ. Sci. Technol.* **50**, 8485–8496 (2016).
136. Soana, E., Balestrini, R., Vincenzi, F., Bartoli, M. & Castaldelli, G. Mitigation of nitrogen pollution in vegetated ditches fed by nitrate-rich spring waters. *Agric. Ecosyst. Environ.* **243**, 74–82 (2017).
137. Vymazal, J. & Březinová, T. D. Removal of nutrients, organics and suspended solids in vegetated agricultural drainage ditch. *Ecol. Eng.* **118**, 97–103 (2018).
138. Baker, B. H., Kröger, R., Brooks, J. P., Smith, R. K. & Prince Czarniecki, J. M. Investigation of denitrifying microbial communities within an agricultural drainage system fitted with low-grade weirs. *Water Res.* **87**, 193–201 (2015).
139. Pierobon, E., Castaldelli, G., Mantovani, S., Vincenzi, F. & Fano, E. A. Nitrogen removal in vegetated and unvegetated drainage ditches impacted by diffuse and point sources of pollution. *CLEAN–Soil Air Water* **41**, 24–31 (2013).
140. Castaldelli, G. et al. Vegetated canals mitigate nitrogen surplus in agricultural watersheds. *Agric. Ecosyst. Environ.* **212**, 253–262 (2015).
141. McPhillips, L. E., Groffman, P. M., Schneider, R. L. & Walter, M. T. Nutrient cycling in grassed roadside ditches and lawns in a suburban watershed. *J. Environ. Qual.* **45**, 1901–1909 (2016).
142. Tatarw, C., Mason, O. U. & Mortazavi, B. Ditching nutrients: roadside drainage networks are hotspots for microbial nitrogen removal. *J. Geophys. Res.: Biogeosci.* **126**, e2020JG006115 (2021).
143. van Liere, L., Janse, J. H. & Arts, G. H. Setting critical nutrient values for ditches using the eutrophication model PCDitch. *Aquat. Ecol.* **41**, 443–449 (2007).
144. Søndergaard, M., Jensen, J. P. & Jeppesen, E. Role of sediment and internal loading of phosphorus in shallow lakes. *Hydrobiologia* **506**, 135–145 (2003).
145. Hill, C. R. & Robinson, J. S. Phosphorus flux from wetland ditch sediments. *Sci. Total Environ.* **437**, 315–322 (2012).
146. McDowell, R. W., Depree, C. & Stenger, R. Likely controls on dissolved reactive phosphorus concentrations in baseflow of an agricultural stream. *J. Soils Sediment.* **20**, 3254–3265 (2020).
147. Trentman, M. T., Tank, J. L., Jones, S. E., McMillian, S. K. & Royer, T. V. Seasonal evaluation of biotic and abiotic factors suggests phosphorus retention in constructed floodplains in three agricultural streams. *Sci. Total Environ.* **729**, 138744 (2020).
148. Kleinman, P. J., Smith, D. R., Bolster, C. H. & Easton, Z. M. Phosphorus fate, management, and modeling in artificially drained systems. *J. Environ. Qual.* **44**, 460–466 (2015).
149. Mhret Gela, S. & Aragaw, T. A. Abundance and characterization of microplastics in main urban ditches across the Bahir Dar City, Ethiopia. *Front. Environ. Sci.* **10**, 35 (2022).
150. McDowell, R. W., Macintosh, K. A. & Depree, C. Linking the uptake of best management practices on dairy farms to catchment water quality improvement over a 20-year period. *Sci. Total Environ.* **895**, 164963 (2023).
151. Pruden, A., Pei, R., Storteboom, H. & Carlson, K. H. Antibiotic resistance genes as emerging contaminants: studies in northern Colorado. *Environ. Sci. Technol.* **40**, 7445–7450 (2006).
152. Gall, H. E., Sassman, S. A., Jenkinson, B., Lee, L. S. & Jafvert, C. T. Comparison of export dynamics of nutrients and animal-borne estrogens from a tile-drained Midwestern agroecosystem. *Water Res.* **72**, 162–173 (2015).
153. de Santiago-Martín, A. et al. Pharmaceuticals and trace metals in the surface water used for crop irrigation: Risk to health or natural attenuation?. *Sci. Total Environ.* **705**, 135825 (2020).
154. Kumwimba, M. N., Zhu, B., Wang, T., Yuan, Z. & Muyembe, D. K. Metal distribution and contamination assessment in drainage ditch water in the main rice/vegetable area of Sichuan hilly basin. *Bull. Environ. Contamin. Toxicol.* **96**, 248–253 (2016).
155. Smiley, P. C. Jr., King, K. W. & Fausey, N. R. Annual and seasonal differences in pesticide mixtures within channelized agricultural headwater streams in central Ohio. *Agric. Ecosyst. Environ.* **193**, 83–95 (2014).
156. Brunet, N. N. & Westbrook, C. J. Wetland drainage in the Canadian prairies: Nutrient, salt and bacteria characteristics. *Agric. Ecosyst. Environ.* **146**, 1–12 (2012).
157. Pan, T. & Miao, T. Contamination of roadside soils by runoff pollutants: A numerical study. *Transport. Geotech.* **2**, 1–9 (2015).
158. Trenouth, W. R. & Gharabaghi, B. Highway runoff quality models for the protection of environmentally sensitive areas. *J. Hydrol.* **542**, 143–155 (2016).
159. Reinikainen, J., Perkola, N., Äystö, L. & Sorvari, J. The occurrence, distribution, and risks of PFAS at AFFF-impacted sites in Finland. *Sci. Total Environ.* **829**, 154237 (2022).
160. Boger, A. R., Ahiablame, L., Mosase, E. & Beck, D. Effectiveness of roadside vegetated filter strips and swales at treating roadway runoff: A tutorial review. *Environ. Sci. Water Res. Technol.* **4**, 478–486 (2018).
161. Moore, M. T. et al. Vegetated ditches for mitigation of contaminants in agricultural runoff. In *The Ecotoxicology of Aquatic Macrophytes* (171–192) (Cham: Springer International Publishing, 2023).
162. Tromp, K., Lima, A. T., Barendregt, A. & Verhoeven, J. T. Retention of heavy metals and poly-aromatic hydrocarbons from road water in a constructed wetland and the effect of de-icing. *J. Hazard. Mater.* **203**, 290–298 (2012).
163. Krause, L. et al. Impacts of historical ditching on peat volume and carbon in northern Minnesota USA peatlands. *J. Environ. Manag.* **296**, 113090 (2021).
164. Evans, C. D., Renou-Wilson, F. & Strack, M. The role of waterborne carbon in the greenhouse gas balance of drained and re-wetted peatlands. *Aquat. Sci.* **78**, 573–590 (2016).
165. Mitchell, G. N. Water quality issues in the British uplands. *Appl. Geogr.* **11**, 201–214 (1991).
166. Roulet, N. T. & Moore, T. R. The effect of forestry drainage practices on the emission of methane from northern peatlands. *Can. J. Res.* **25**, 491–499 (1995).
167. Deng, O. et al. CH₄ and CO₂ emissions in water networks of rice cultivation regions. *Environ. Res.* **218**, 115041 (2023).
168. IPCC. Refinement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories. Volume 4: Agriculture, forestry and other land use (AFOLU). Chapter 7: Wetlands. Lovelock, C.E., Evans, C., Barros, N., Prairie, Y., Alm, J., Bastviken, D., Beaulieu, J.J., Garneau, M., Harby, A., Harrison, J., Pare, D., Raadal, H.L.,

- Sherman, B., Zhang, C., Ogle, S.M., Grinham, A., Deemer, B., Aurelio dos Santos, M., Kosten, S., Peacock, M., Li, Z. & Stepanenko, V. Intergovernmental Panel on Climate Change (IPCC, Switzerland 2019).
169. Iseyemi, O. et al. Characterizing organic carbon storage in experimental agricultural ditch systems in Northeast Arkansas. *Soil Sci. Soc. Am. J.* **83**, 751–760 (2019).
170. Clifford, C. C. & Heffernan, J. B. North Carolina Coastal Plain ditch types support distinct hydrophytic communities. *Wetlands* **43**, 56 (2023).
171. Needelman, B. A., Ruppert, D. E. & Vaughan, R. E. The role of ditch soil formation and redox biogeochemistry in mitigating nutrient and pollutant losses from agriculture. *J. Soil Water Conserv.* **62**, 207–215 (2007).
172. Peacock, M., Granath, G., Wallin, M. B., Högbom, L. & Futter, M. N. Significant emissions from forest drainage ditches—An unaccounted term in Anthropogenic greenhouse gas inventories? *J. Geophys. Res.: Biogeosci.* **126**, e2021JG006478 (2021).
173. Hendriks, L. et al. Drainage ditches are year-round greenhouse gas hotlines in temperate peat landscapes. *Freshw. Biol.* **69**, 143–156 (2024).
174. Tanneberger, F. et al. Towards net zero CO₂ in 2050: An emission reduction pathway for organic soils in Germany. *Mires Peat* **27**, 17pp (2021).
175. Bowen, J. C., Wahyudjo, P. J., Anshari, G. Z., Aluwihare, L. I. & Hoyt, A. M. Canal networks regulate aquatic losses of carbon from degraded tropical peatlands. *Nat. Geosci.* **17**, 213–218 (2024).
176. Evans, C. & Taillardat, P. Light on dark waters. *Nat. Geosci.* **17**, 174–175 (2024).
177. Silverthorn, T. et al. The importance of ditches and canals in global inland water CO₂ and N₂O budgets. *Glob. Chang. Biol.* **31**, e70079 (2025).
178. Clagnan, E. et al. Investigating “net” provenance, N source, transformation and fate within hydrologically isolated grassland plots. *Agric. Water Manag.* **203**, 1–8 (2018).
179. Kröger, R., Holland, M. M., Moore, M. T. & Cooper, C. M. Hydrological variability and agricultural drainage ditch inorganic nitrogen reduction capacity. *J. Environ. Qual.* **36**, 1646–1652 (2007).
180. Reay, D. S., Smith, K. A. & Edwards, A. C. Nitrous oxide emission from agricultural drainage waters. *Glob. Change Biol.* **9**, 195–203 (2003).
181. Nevison, C. Review of the IPCC methodology for estimating nitrous oxide emissions associated with agricultural leaching and runoff. *Chemosphere-Glob. Change Sci.* **2**, 493–500 (2000).
182. Webb, J. R., Clough, T. J. & Quayle, W. C. A review of indirect N₂O emission factors from artificial agricultural waters. *Environ. Res. Lett.* **16**, 043005 (2021).
183. Beaulieu, J. J. et al. Nitrous oxide emission from denitrification in stream and river networks. *PNAS* **108**, 214–219 (2011).
184. Armitage, P. D., Szoszkiewicz, K., Blackburn, J. H. & Nesbitt, I. Ditch communities: a major contributor to floodplain biodiversity. *Aquat. Conserv. Mar. Freshw. Ecosyst.* **13**, 165–185 (2003).
185. Painter, D. Macroinvertebrate distributions and the conservation value of aquatic Coleoptera, Mollusca and Odonata in the ditches of traditionally managed and grazing fen at Wicken Fen, UK. *J. Appl. Ecol.* **36**, 33–48 (1999).
186. Law, A. et al. Repeatable patterns in the distribution of freshwater biodiversity indicators across contrasting landscapes. *Landsc. Ecol.* **39**, 195 (2024).
187. Hill, M. J., Chadd, R. P., Morris, N., Swaine, J. D. & Wood, P. J. Aquatic macroinvertebrate biodiversity associated with artificial agricultural drainage ditches. *Hydrobiologia* **776**, 249–260 (2016).
188. King, S., Elphick, C. S., Guadagnin, D., Taft, O. & Amano, T. Effects of landscape features on waterbird use of rice fields. *Waterbirds* **33**, 151–159 (2010).
189. Marja, R. & Herzon, I. The importance of drainage ditches for farmland birds in agricultural landscapes in the Baltic countries: does field type matter?. *Ornis Fennica* **89**, 170–181 (2012).
190. Johnson, B. A., Barrett, K., Homyack, J. A. & Baldwin, R. F. Anuran occupancy and breeding site use of aquatic systems in a managed pine landscape. *Ecol. Manag.* **368**, 45–54 (2016).
191. Hinojosa-Garro, D., Mason, C. F. & Underwood, G. J. Macrophyte assemblages in ditches of coastal marshes in relation to land-use, salinity and water quality. *Fundam. Appl. Limnol.* **172**, 325 (2008).
192. Meier, M., Gerlach, R., Schirmel, J. & Buhk, C. Plant diversity in a water-meadow landscape: the role of irrigation ditches. *Plant Ecol.* **218**, 971–981 (2017).
193. Ryšavá-Nováková, M., Ondračková, M. & Jurajda, P. The importance of surrogate habitats in lowland river floodplains for fish community composition. *Fish. Manag. Ecol.* **16**, 468–477 (2009).
194. Rosenvald, R., Järvekülg, R. & Lohmus, A. Fish assemblages in forest drainage ditches: Degraded small streams or novel habitats?. *Limnologica* **46**, 37–44 (2014).
195. Sun, J. et al. Regional macrophyte diversity is shaped by accumulative effects across waterbody types in southern China. *Aquat. Bot.* **176**, 103468 (2022).
196. Van Geert, A., Van Rossum, F. & Triest, L. Do linear landscape elements in farmland act as biological corridors for pollen dispersal?. *J. Ecol.* **98**, 178–187 (2010).
197. O'Bryan, C. J., Homyack, J. A., Baldwin, R. F., Kanno, Y. & Harrison, A. L. Novel habitat use supports population maintenance in a reconfigured landscape. *Ecosphere* **7**, e01228 (2016).
198. Teurlincx, S., Verhofstad, M. J., Bakker, E. S. & Declerck, S. A. Managing successional stage heterogeneity to maximize landscape-wide biodiversity of aquatic vegetation in ditch networks. *Front. Plant Sci.* **9**, 1013 (2018).
199. Gething, K. J., Sykes, T., Biondi, G., Macadam, C. & Stubbington, R. Ditching misconceptions: rare temporary stream specialists in artificial habitats. *FBA News Freshw. Biol. Assoc. Newsl.* **83**, 14–18 (2021).
200. Lohmus, A., Remm, L. & Rannap, R. Just a ditch in forest? Reconsidering draining in the context of sustainable forest management. *BioScience* **65**, 1066–1076 (2015).
201. Vaikre, M., Remm, L. & Rannap, R. Forest ditch maintenance impoverishes the fauna of aquatic invertebrates: opportunities for mitigation. *J. Environ. Manag.* **274**, 111188 (2020).
202. Perrone, S. M., Deutsch, C., Bilenca, D. N. & Agostini, M. G. Artificial aquatic habitats impoverish amphibian diversity in agricultural landscapes of central Argentina. *Aquat. Conserv. Mar. Freshw. Ecosyst.* **32**, 591–604 (2022).
203. Clarke, S. J. Conserving freshwater biodiversity: the value, status and management of high quality ditch systems. *J. Nat. Conserv.* **24**, 93–100 (2015).
204. Gardner, R. C. & Finlayson, C. *Ramsar Convention on Wetlands. Global Wetland Outlook: State of the World's Wetlands and their Services to People* (Gland, Switzerland: Ramsar Convention Secretariat, 2018).
205. Langheinrich, U., Tischew, S., Gersberg, R. M. & Lüderitz, V. Ditches and canals in management of fens: opportunity or risk? A case study in the Drömling Natural Park, Germany. *Wetl. Ecol. Manag.* **12**, 429–445 (2004).
206. Suren, A. M., Lambert, P. & Sorrell, B. K. The impact of hydrological restoration on benthic aquatic invertebrate communities in a New Zealand wetland. *Restor. Ecol.* **19**, 747–757 (2011).
207. Friday, L. E. & Harley B. (eds) *Checklist of the Flora and Fauna of Wicken Fen* (Harley Books, England 2000).
208. Watson, A. M. & Ormerod, S. J. The microdistribution of three uncommon freshwater gastropods in the drainage ditches of British grazing marshes. *Aquat. Conserv. Mar. Freshw. Ecosyst.* **14**, 221–236 (2004).

209. Michna, P. 552. *Senecio paludosus*: Compositae. *Curtis Botanical Mag.* **23**, 69–76 (2006).
210. Mazerolle, M. J. Drainage ditches facilitate frog movements in a hostile landscape. *Landsc. Ecol.* **20**, 579–590 (2005).
211. Favre-Bac, L. et al. Connectivity and propagule sources composition drive ditch plant metacommunity structure. *Acta Oecol.* **61**, 57–64 (2014).
212. Salazar, R. D., Montgomery, R. A., Thresher, S. E. & Macdonald, D. W. Mapping the relative probability of common toad occurrence in terrestrial lowland farm habitat in the United Kingdom. *PLoS One* **11**, e0148269 (2016).
213. Bouchard, E. H. et al. Landscape factors driving the spread of the invasive grass, *Hymenachne amplexicaulis*, among wetlands in a Florida subtropical grazing land. *Invasive Plant Sci. Manag.* **13**, 155–162 (2020).
214. Goeller, B. C., Sukias, J. P., Woodward, S. J. & Clarkson, B. R. Dual purpose surface flow constructed treatment wetlands support native biodiversity in intensified agricultural landscapes. *Water* **15**, 2526 (2023).
215. Vindigni, M. A., Morris, A. D., Miller, D. A. & Kalcounis-Rueppell, M. C. Use of modified water sources by bats in a managed pine landscape. *Ecol. Manag.* **258**, 2056–2061 (2009).
216. Raitif, J., Plantegenest, M. & Roussel, J. M. From stream to land: Ecosystem services provided by stream insects to agriculture. *Agric. Ecosyst. Environ.* **270**, 32–40 (2019).
217. Wen, Y. et al. Raising the groundwater table in the non-growing season can reduce greenhouse gas emissions and maintain crop productivity in cultivated fen peats. *J. Clean. Prod.* **262**, 121179 (2020).
218. Zak, D. et al. An assessment of the multifunctionality of integrated buffer zones in Northwestern Europe. *J. Environ. Qual.* **48**, 362–375 (2019).
219. Olson, D. H. & Burnett, K. M. Geometry of forest landscape connectivity: pathways for persistence. Density Management in the 21st Century: West Side Story. General Technical Report PNW-GTR-880. US Department of Agriculture, Forest Service, Pacific Northwest Research Station. Portland, Oregon, USA, 220–238 (2013).
220. Churchman, C. W. Guest editorial: Wicked problems. *Manag. Sci.* **14**, B141–B142 (1967).
221. Palta, M. M., Grimm, N. B. & Groffman, P. M. Accidental urban wetlands: ecosystem functions in unexpected places. *Front. Ecol. Environ.* **15**, 248–256 (2017).
222. Borja, A. & Elliott, M. What does ‘good ecological potential’ mean, within the European Water Framework Directive?. *Mar. Pollut. Bull.* **54**, 1559–1564 (2007).
223. Sayer, C. D. Conservation of aquatic landscapes: ponds, lakes, and rivers as integrated systems. *Wiley Interdiscip. Rev.: Water* **1**, 573–585 (2014).
224. Loreau, M., Mouquet, N. & Holt, R. D. Meta-ecosystems: a theoretical framework for a spatial ecosystem ecology. *Ecol. Lett.* **6**, 673–679 (2003).
225. Hambäck, P. A. et al. Tradeoffs and synergies in wetland multifunctionality: A scaling issue. *Sci. Total Environ.* **862**, 160746 (2022).
226. Steele, M. K. et al. Convergent surface water distributions in US cities. *Ecosystems* **17**, 685–697 (2014).
227. Boothby, J. Pond conservation: towards a delineation of pondscape. *Aquat. Conserv.: Mar. Freshw. Ecosyst.* **7**, 127–132 (1997).
228. Allan, J. D. Landscapes and riverscapes: the influence of land use on stream ecosystems. *Annu. Rev. Ecol. Evol. Syst.* **35**, 257–284 (2004).
229. Temmink, R. J. et al. Wetscapes: Restoring and maintaining peatland landscapes for sustainable futures. *Ambio* **52**, 1519–1528 (2023).
230. Stratigos, M. J. What wetland are we protecting and restoring? Quantifying the human creation of protected areas in Scotland. *Journal of Wetland Archaeology*, 1–17 (2022).
231. Connolly, J. & Holden, N. M. Detecting peatland drains with object based image analysis and Geoeye-1 imagery. *Carbon balance Manag.* **12**, 1–13 (2017).
232. Dadap, N. C. et al. Drainage canals in Southeast Asian peatlands increase carbon emissions. *AGU Adv.* **2**, e2020AV000321 (2021).
233. Robb, C., Pickard, A., Williamson, J. L., Fitch, A. & Evans, C. Peat drainage ditch mapping from aerial imagery using a convolutional neural network. *Remote Sens.* **15**, 499 (2023).
234. Lidberg, W. et al. Mapping drainage ditches in forested landscapes using deep learning and aerial laser scanning. *J. Irrig. Drain. Eng.* **149**, 04022051 (2023).
235. Habib, W., Cresson, R., McGuinness, K. & Connolly, J. Mapping artificial drainage on Peatlands – A national scale assessment of Irish raised bogs using sub-meter aerial imagery and deep learning methods. *Remote Sens. Ecol. Conserv.* **10**, 551–562 (2024).
236. Fritz, K., Cid, N. & Autrey, B. Governance, legislation, and protection of intermittent rivers and ephemeral streams. In: *Intermittent rivers and ephemeral streams* (477–507). Academic Press (2017).
237. Williams, P. et al. Comparative biodiversity of rivers, streams, ditches and ponds in an agricultural landscape in Southern England. *Biol. Conserv.* **115**, 329–341 (2004).
238. Brown, C. D. et al. Morphological and physico-chemical properties of British aquatic habitats potentially exposed to pesticides. *Agric. Ecosyst. Environ.* **113**, 307–319 (2006).
239. Davies, B. et al. Comparative biodiversity of aquatic habitats in the European agricultural landscape. *Agric. Ecosyst. Environ.* **125**, 1–8 (2008).
240. Davies, B., Biggs, J., Williams, P. & Thompson, S. Making agricultural landscapes more sustainable for freshwater biodiversity: a case study from southern England. *Aquat. Conserv.: Mar. Freshw. Ecosyst.* **19**, 439–447 (2009).
241. Shaw, R. F., Johnson, P. J., Macdonald, D. W. & Feber, R. E. Enhancing the biodiversity of ditches in intensively managed UK farmland. *PLoS One* **10**, e0138306 (2015).
242. Bubíková, K. & Hrivnák, R. Comparative macrophyte diversity of waterbodies in the Central European landscape. *Wetlands* **38**, 451–459 (2018).
243. Nakano, M. & Morii, K. Factors affecting the abundance of a clam (genus *Corbicula*) and distribution overlap between the clam and unionids in agricultural ditches. *Jpn. J. Environ. Entomol. Zool.* **30**, 1–8 (2019).
244. Williams, P. et al. Nature based measures increase freshwater biodiversity in agricultural catchments. *Biol. Conserv.* **244**, 108515 (2020).
245. Gething, K. J. & Little, S. The importance of artificial drains for macroinvertebrate biodiversity in reclaimed agricultural landscapes. *Hydrobiologia* **847**, 3129–3138 (2020).
246. Verdonschot, R. C., Keizer-Vlek, H. E. & Verdonschot, P. F. Development of a multimetric index based on macroinvertebrates for drainage ditch networks in agricultural areas. *Ecol. Indic.* **13**, 232–242 (2012).
247. Cooper, C. M., Moore, M. T., Bennett, E. R., Smith, S. Jr & Farris, J. L. Alternative environmental benefits of agricultural drainage ditches. *Int. Ver. f. ü. theoretische und Angew. Limnologie: Verhandlungen* **28**, 1678–1682 (2002).
248. Fu, D. et al. Nutrient mitigation capacity of agricultural drainage ditches in Tai lake basin. *Ecol. Eng.* **71**, 101–107 (2014).
249. Shore, M., Jordan, P., Mellander, P. E., Kelly-Quinn, M. & Melland, A. R. An agricultural drainage channel classification system for phosphorus management. *Agric. Ecosyst. Environ.* **199**, 207–215 (2015).

250. Pierce, S. C., Kröger, R. & Pezeshki, R. Managing artificially drained low-gradient agricultural headwaters for enhanced ecosystem functions. *Biology* **1**, 794–856 (2012).
251. Ahmed, F., Gulliver, J. S. & Nieber, J. L. Field infiltration measurements in grassed roadside drainage ditches: Spatial and temporal variability. *J. Hydrol.* **530**, 604–611 (2015).
252. Nsenga Kumwimba, M. et al. Assessing nutrient, biomass, and sediment transport of drainage ditches in the Three Gorges Reservoir Area. *CLEAN–Soil, Air, Water*, **45** (2017).
253. European Protection Agency. Ditch. Available from: <https://www.eea.europa.eu/help/glossary/gemet-environmental-thesaurus/ditch> [Last accessed: 21/4/2023] 2023..

Acknowledgements

We thank others who attended the “Raising the Profile of Ditch Research” workshop or provided comments on the draft, but did not want to contribute as named authors. Funding: Mike Peacock: Natural Environment Research Council [NE/X018423/1], Formas projects LEAF-PAD [2020-00950] and PUDDLE-JUMP [2022-02138], Naturvårdsverket project WetKit Hydro-ES [802-0083-19]. Joachim Audet: Independent Research Fund Denmark project DrivNOS grant nr. 0217-00021B. Laurie Friday and Hamidreza Rahimi: Natural Environment Research Council Changing the Environment Programme [NE/W00495X/1]. Ivan Arismendi and Guillermo Giannico received support from the House Bill 2437, Chapter 699, Oregon Legislative Assembly, 2019. We thank Richard McDowell for providing references to research on Australasian ditches, and two anonymous reviewers for their constructive comments that helped to improve the paper.

Author contributions

Chelsea Clifford initiated the informal group of ditch researchers who wrote this paper, expanded by Mike Peacock. Chelsea Clifford and Mike Peacock conceived the paper and coordinated authors. Mike Peacock acquired funding for a two-day NERC-funded workshop “Raising the Profile of Ditch Research” during which the paper was planned by many of its authors. Chelsea Clifford and Mike Peacock led the paper, and the following authors coordinated/led the writing of individual sections: Chelsea Clifford (Introduction), Michael Stratigos (Human) (with support from Nejem Raheem), Magdalena Bieroza (Physical), Stewart Clarke (Flora and Fauna) (with support from Matt Hill and Paul Wood), Amy Pickard (Biogeochemistry) (with support from Corianne Tatarwi), Mike Peacock (Synthesis & Conclusions). Matt Hill compiled ditch/channel definitions in Table 1. John Connolly coined the phrase “lines in the landscape”. Mans Schepers contributed Fig. 2. Ivan Arismendi, Joachim Audet, Daniel Aviles, Jordanna N. Bergman, Anthony G. Brown, Rachel Eleanor Burns, John Connolly, Sarah Cook, Julie Crabot, Wyatt F. Cross, Joshua F. Dean, Chris D. Evans, Owen Fenton, Laurie Friday, Kieran J. Gething, Guillermo Giannico, Wahaj Habib, Eliza Maher Hasselquist, Nathaniel M. Heili, Judith van der Knaap, Sarian Kosten, Alan Law, Gea H. van der Lee, Kate L. Mathers, John E. Morgan, Hamidreza Rahimi, Carl D.

Sayer, Mans Schepers, Rosalind F. Shaw, Peter C. Smiley Jr., Shannon L. Speir, Jeffrey S. Strock, Quinten Struik, Jennifer L. Tank, Hao Wang, Jackie R. Webb, Alex J. Webster, Zhifeng Yan, and Peta Zivec contributed ideas and text, read and commented on drafts, and approved the final manuscript.

Funding

Open access funding provided by Swedish University of Agricultural Sciences

Competing interests

Joshua Dean is an Editorial Board Member for Communications Earth & Environment, but was not involved in the editorial review of, nor the decision to publish this article. The authors declare no competing interests.

Additional information

Supplementary information The online version contains supplementary material available at <https://doi.org/10.1038/s43247-025-02699-y>.

Correspondence and requests for materials should be addressed to Chelsea Clifford or Mike Peacock.

Peer review information *Communications Earth & Environment* thanks Brandon C. Goeller and the other, anonymous, reviewer(s) for their contribution to the peer review of this work. Primary Handling Editors: Martina Grecequet A peer review file is available.

Reprints and permissions information is available at <http://www.nature.com/reprints>

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2025

¹Virginia Institute of Marine Science, College of William and Mary, Gloucester Point, VA, USA. ²Department of Soil and Environment, Swedish University of Agricultural Sciences, Uppsala, Sweden. ³National Trust, Heelis, Swindon, UK. ⁴UK Centre for Ecology & Hydrology, Bush Estate, Penicuik, Scotland. ⁵Interdisciplinary Institute and Department of Archaeology, St Mary's Building, Elphinstone Road, University of Aberdeen, Aberdeen, UK. ⁶Department of Agriculture and Environment, Harper Adams University, Newport, UK. ⁷Department of Marketing Communication, Emerson College, Boston, MA, USA. ⁸Department of Environmental Science, Rowan University, Glassboro, NJ, USA. ⁹Geography and Environment, Loughborough University, Loughborough, Leicestershire, UK. ¹⁰Department of Fisheries, Wildlife, and Conservation Sciences, Oregon State University, Corvallis, OR, USA. ¹¹Department of Ecoscience, Aarhus University, Aarhus, Denmark. ¹²Universidad Mayor de San Simon, Laboratorio de Hidraulica, Cochabamba, Bolivia. ¹³Department of Biology, Carleton University, Ottawa, ON, Canada. ¹⁴Botany Section, Tromsø Museum, Arctic University of Norway, Tromsø, Norway. ¹⁵Geography and Environment, University of Southampton, Southampton, UK. ¹⁶Department of Geosciences and Natural Resource Management, University of Copenhagen, Copenhagen, Denmark. ¹⁷Discipline of Geography, School of Natural Sciences, Trinity College Dublin, Dublin, Ireland. ¹⁸Department of Life Sciences, University of Warwick, Coventry, UK. ¹⁹FEHM-Lab (Freshwater Ecology, Hydrology and Management), Institute of Environmental Assessment and Water Research (IDAEA), CSIC, Barcelona, Spain. ²⁰Department of Ecology, Montana State University, Bozeman, MT, USA. ²¹School of Geographical Sciences, University of Bristol, Bristol, UK. ²²UK Centre for Ecology and Hydrology, Bangor, UK. ²³Environment, Soils and Land Use Department, Teagasc Johnstown Castle Research Centre, Wexford, Ireland. ²⁴Centre for Landscape Regeneration and Department of Plant Sciences, University of Cambridge, Cambridge, UK. ²⁵School of Science and Technology, Nottingham Trent University, Nottingham, UK. ²⁶Department of Forest Ecology and Management, Swedish

University of Agricultural Sciences, Umeå, Sweden. ²⁷Department of Ecology, Radboud Institute for Biological and Environmental Sciences, Radboud University, Nijmegen, the Netherlands. ²⁸Biological and Environmental Sciences, University of Stirling, Stirling, UK. ²⁹Wageningen Environmental Research, Wageningen UR, Wageningen, the Netherlands. ³⁰Pond Restoration Research Group, Department of Geography, University College London, London, UK. ³¹Groningen Institute of Archaeology, University of Groningen, Groningen, The Netherlands. ³²Environment and Sustainability Institute, University of Exeter, Penryn Campus, Penryn, Cornwall, UK. ³³USDA Agricultural Research Service, Columbus, OH, USA. ³⁴Department of Crop, Soil, and Environmental Science, University of Arkansas, Fayetteville, AR, USA. ³⁵Department of Soil, Water, and Climate, University of Minnesota, St. Paul, MN, USA. ³⁶Southwest Research and Outreach Center, University of Minnesota, Lamberton, MN, USA. ³⁷Department of Biological Sciences, University of Notre Dame, Notre Dame, IN, USA. ³⁸School of Agriculture & Environmental Science, University of Southern Queensland, Toowoomba, QLD, Australia. ³⁹Centre for Sustainable Agricultural Systems, University of Southern Queensland, Toowoomba, QLD, Australia. ⁴⁰Department of Biology, University of New Mexico, Albuquerque, NM, USA. ⁴¹Institute of Surface-Earth System Science, School of Earth System Science, Tianjin University, Tianjin, China. ⁴²Department of Aquatic Sciences and Assessment, Swedish University of Agricultural Sciences, Uppsala, Sweden. ⁴³Department of Geography and Planning, School of Environmental Sciences, University of Liverpool, Liverpool, UK. ⁴⁴These authors contributed equally: Chelsea Clifford, Mike Peacock. ✉ e-mail: ccclifford@vims.edu; chelseaclifford@gmail.com; m.peacock@liverpool.ac.uk; michael.peacock@slu.se